

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 May 2001 (31.05.2001)

PCT

(10) International Publication Number
WO 01/38397 A1

- (51) International Patent Classification⁷: C07K 14/78, 7/08
- (21) International Application Number: PCT/US00/32105
- (22) International Filing Date:
22 November 2000 (22.11.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/447,099 22 November 1999 (22.11.1999) US
09/702,649 31 October 2000 (31.10.2000) US
- (71) Applicant: ABBOTT LABORATORIES [US/US];
Dept. 377/AP6D-2, 100 Abbott Park Road, Abbott Park,
IL 60064-6050 (US).
- (72) Inventors: HAVIV, Fortuna; 1125 Oxford Road,
Deerfield, IL 60015 (US). HENKIN, Jack; 1370 Lincoln
Avenue South, Highland Park, IL 60035 (US). BRADLEY,
Michael, F.; 2591 North Augusta Drive, Wadsworth, IL
60083 (US). KALVIN, Douglas, M.; 1201 Lockwood
Drive, Buffalo Grove, IL 60089 (US).
- (54) Agents: DONNER, B., Gregory et al.; Dept.
377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL
60064-6050 (US).
- (81) Designated States (*national*): AL, AG, AL, AM, AI, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/38397 A1

(54) Title: N-ALKYLATED PEPTIDES HAVING ANTIANGIOGENIC ACTIVITY

(57) Abstract: N-Alkylated peptides of the formula (I) Xaa₁-Xaa₂-Xaa₃-Xaa₄-Xaa₅-Xaa₆-Xaa₇-Xaa₈-Xaa₉-Xaa₁₀-Xaa₁₁, are useful for inhibiting angiogenesis. Also disclosed are angiogenesis-inhibiting compositions and methods of inhibiting angiogenesis in a mammal.

5 **N-ALKYLATED PEPTIDES HAVING ANTIANGIOGENIC ACTIVITY**

Technical Field

 The invention relates to novel compounds having activity useful for treating conditions which arise or are exacerbated by angiogenesis, pharmaceutical compositions
10 comprising the compounds, methods of treatment using the compounds, and methods of inhibiting angiogenesis.

Background of the Invention

 Angiogenesis is the fundamental process by which new blood vessels are formed
15 and is essential to a variety of normal body activities (such as reproduction, development and wound repair). Although the process is not completely understood, it is believed to involve a complex interplay of molecules which both stimulate and inhibit the growth of endothelial cells, the primary cells of the capillary blood vessels. Under normal conditions these molecules appear to maintain the microvasculature in a quiescent state (i.e., one of
20 no capillary growth) for prolonged periods that may last for weeks, or in some cases, decades. However, when necessary, such as during wound repair, these same cells can undergo rapid proliferation and turnover within as little as five days (Folkman, J. and Shing, Y., *The Journal of Biological Chemistry*, **267**(16): 10931-10934, and Folkman, J. and Klagsbrun, M., *Science*, **235**: 442-447 (1987)).

25 Although angiogenesis is a highly regulated process under normal conditions, many diseases (characterized as "angiogenic diseases") are driven by persistent unregulated angiogenesis. Otherwise stated, unregulated angiogenesis may either cause a particular disease directly or exacerbate an existing pathological condition. For example, ocular neovascularization has been implicated as the most common cause of blindness. In
30 certain existing conditions such as arthritis, newly formed capillary blood vessels invade the joints and destroy cartilage. In diabetes, new capillaries formed in the retina invade the vitreous, bleed, and cause blindness. Growth and metastasis of solid tumors are also angiogenesis-dependent (Folkman, J., *Cancer Research*, **46**: 467-473 (1986), Folkman, J., *Journal of the National Cancer Institute*, **82**: 4-6 (1989)). It has been shown, for example,
35 that tumors which enlarge to greater than 2 mm, must obtain their own blood supply and do so by inducing the growth of new capillary blood vessels. Once these new blood

vessels become embedded in the tumor, they provide a means for tumor cells to enter the circulation and metastasize to distant sites, such as the liver, the lung, and the bones (Weidner, N., et. al., *The New England Journal of Medicine*, **324**(1): 1-8 (1991)).


Several angiogenesis inhibitors are currently under development for use in treating angiogenic diseases (Gasparini, G. and Harris, A.L., *J Clin Oncol* **13**(3): 765-782, (1995)). A number of disadvantages have been associated with many of these compounds. A potent angiogenesis inhibitor, for example suramin, can cause severe systemic toxicity in humans at doses required to reach antitumor activity. Other compounds, such as retinoids, interferons, and antiestrogens are safe for human use, but have only a weak anti-angiogenic effect.

A new class of compounds having particularly effective *in vitro* and *in vivo* angiogenesis inhibiting properties, as well as a promising toxicity profile, has been described in commonly-owned U.S. Patent Application No. 09/316,888, filed May 21, 1999. Novel changes in position 3 of the new angiogenesis inhibitors is described in copending provisional U.S. Patent Application Ser. No. 60/166,791, filed November 22, 1999. Although the compounds have demonstrated satisfactory enzymatic stability and bioavailability, it would be desirable to prepare analogs of the antiangiogenic peptides having enhanced stability against *in vivo* enzymatic cleavage, improved pharmacokinetics, increased water solubility, and potentially better oral bioavailability.

Summary of the Invention

The present invention relates to a novel class of compounds having angiogenesis-inhibiting properties. The invention provides nona- and decapeptides wherein the nitrogen atom of at least one of the amide bonds of an amino acid residue in positions 2 through 9 of the peptide is N-alkylated. Compounds of the invention exhibit enhanced metabolic stability, improved pharmacokinetics, increased water solubility, and potentially better oral bioavailability.

In one aspect, the present invention provides a compound of formula (I)



or a pharmaceutically acceptable salt, prodrug, ester, or solvate thereof, wherein

at least one amide bond of an amino acid residue represented by Xaa₃, Xaa₄, Xaa₅, Xaa₆, Xaa₇, Xaa₈, Xaa₉, and Xaa₁₀ is N-alkylated;

Xaa₁ is absent or Xaa₁ is selected from the group consisting of hydrogen, N-methylprolyl, and an acyl group, wherein the acyl group is selected from the group consisting of

5 R¹-(CH₂)_n-C(O)-, wherein n is an integer from 0 to 8 and R¹ is selected from the group consisting of N-acetylamino, alkoxy, alkyl, aryl, carboxy, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy; and
 R²-CH₂CH₂-O-(CH₂CH₂O)_p-CH₂-C(O)-, wherein p is an integer from 1 to 8 and R² is selected from the group consisting of hydrogen, N-acetylamino, and alkyl;

10 provided that Xaa₁ is absent only when Xaa₂ is N-(R³)-prolyl;

Xaa₂ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-glycyl, N-(R³)-norvalyl, and N-(R³)-prolyl, wherein R³ is C₁-C₅-alkyl; or Xaa₂ is an N-unalkylated amino acid selected from the group consisting of

15 β-alanyl,
 D-alanyl,
 4-aminobutyryl,
 (1R,3S)-1-aminocyclopentane-3-carbonyl,
 (1S,3R)-1-aminocyclopentane-3-carbonyl,
 20 (1R,4S)-1-aminocyclopent-2-ene-4-carbonyl,
 (1S,4R)-1-aminocyclopent-2-ene-4-carbonyl,
 asparaginy,
 3-(4-chlorophenyl)alanyl,
 3-(4-cyanophenyl)alanyl,
 25 glutaminyl,
 glutamyl,
 glycyl,
 4-hydroxyprolyl,
 3-(4-methylphenyl)alanyl,
 30 prolyl,
 seryl, and
 threonyl;

35 Xaa₃ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-glycyl, N-(R³)-leucyl, and N-(R³)-phenylalanyl, wherein R³ is as

defined above; or Xaa₃ is an N-unalkylated amino acid selected from the group consisting of

alanyl,
(1S,3R)-1-aminocyclopentane-3-carbonyl,
5 (1S,4R)-1-aminocyclopent-2-ene-4-carbonyl,
asparaginyl,
aspartyl,
3-(3-cyanophenyl)alanyl,
3-(4-cyanophenyl)alanyl,
10 glutaminyl,
glycyl,
leucyl,
lysyl(N-epsilon-acetyl),
3-(4-methylphenyl)alanyl,
15 norvalyl,
prolyl, and
phenylalanyl;

Xaa₄ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
20 alanyl, N-(R³)-glycyl, N-(R³)-homophenylalanyl, N-(R³)-isoleucyl, N-(R³)-leucyl,
N-(R³)-norvalyl, N-(R³)-phenylalanyl, N-(R³)-D-phenylalanyl, N-(R³)-seryl, N-
(R³)-tyrosyl, N-(R³)-valyl, and N-(R³)-D-valyl, wherein R³ is as defined above; or
Xaa₄ is an N-unalkylated amino acid selected from the group consisting of

alanyl,
25 alloisoleucyl,
allylglycyl,
2-aminobutyryl,
(1R,4S)-aminocyclopent-2-ene-4-carbonyl,
asparaginyl,
30 aspartyl,
3-[2-(5-bromothieryl)]alanyl,
3-(3-chlorophenyl)alanyl,
3-(4-chlorophenyl)alanyl,
3-(3-cyanophenyl)alanyl,
35 cyclohexylalanyl,
3-(3,4-dimethoxyphenyl)alanyl,

5 3-(3-fluorophenyl)alanyl,
3-(4-fluorophenyl)alanyl,
glutaminyl,
glycyl,
histidyl,
homophenylalanyl,
homoseryl,
isoleucyl,
leucyl,
10 lysyl(N-epsilon-acetyl),
methionyl,
methionyl(sulfone),
3-(4-methylphenyl)alanyl,
3-(naphth-1-yl)alanyl,
15 3-(naphth-2-yl)alanyl,
norornithyl,
norvalyl,
phenylalanyl,
phenylglycyl,
20 prolyl,
3-(3-pyridyl)alanyl,
3-(4-thiazolyl)alanyl,
3-(2-thienyl)alanyl,
seryl,
25 seryl(O-benzyl),
styrylalanyl,
tryptyl,
tyrosyl,
valyl, and
30 D-valyl;

Xaa_i is an N-alkylated amino acid selected from the group consisting of N-(R³)-D-homophenylalanyl, N-(R³)-D-isoleucyl, N-(R³)-D-leucyl, and N-(R³)-D-phenylalanyl, wherein R³ is as defined above; or Xaa_i is an N-unalkylated amino
35 acid selected from the group consisting of
D-alanyl,

alloisoleucyl,
D-alloisoleucyl,
D-2-aminobutyryl,
D-3-(4-aminophenyl)alanyl,
5 D-asparaginyI,
D-3-(3-benzothienyl)alanyl,
D-*t*-butylglycyl,
D-(chlorophenyl)alanyl,
D-citrullyl,
10 D-3-(3-cyanophenyl)alanyl,
D-cyclohexylalanyl,
cyclohexylglycyl,
D-cysteinyI(S-acetamidomethyl),
D-cysteinyI(S-*t*-butyl),
15 D-3-(3,4-difluorophenyl)alanyl,
D-(3,4-dimethoxyphenyl)alanyl,
D-glutaminyI,
glycyl,
D-homophenylalanyl,
20 D-homoseryl,
isoleucyl,
D-isoleucyl,
D-leucyl,
D-lysyl(N-epsilon-nicotinyI),
25 D-lysyl,
D-methionyl,
D-3-(4-methylphenyl)alanyl,
D-3-(naphth-1-yl)alanyl,
D-3-(naphth-2-yl)alanyl,
30 D-3-(4-nitrophenyl)alanyl,
D-norleucyl,
D-ornithyl,
D-penicillaminyI(S-acetamidomethyl),
D-penicillaminyI(S-benzyl),
35 D-penicillaminyI(S-methyl),
D-penicillaminyI,

D-3-(pentafluorophenyl)alanyl,
 D-phenylalanyl,
 D-prolyl,
 D-seryl(O-benzyl),
 5 D-seryl,
 D-(2-thienyl)alanyl,
 D-threonyl(O-benzyl),
 D-threonyl,
 D-3-(3-trifluoromethylphenyl)alanyl,
 10 D-(3,4,5-trifluorophenyl)alanyl,
 D-tryptyl,
 D-tyrosyl(O-ethyl),
 D-tyrosyl, and
 D-valyl;

15

Xaa₆ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
 aspartyl, N-(R³)-glutamyl, N-(R³)-glycyl, N-(R³)-seryl, N-(R³)-threonyl, N-(R³)-
 threonyl(O-benzyl), and N-(R³)-tyrosyl, wherein R³ is as defined above; or Xaa₆ is
 an N-unalkylated amino acid selected from the group consisting of

20 alanyl,
 allothreonyl,
 D-allothreonyl,
 allylglycyl,
 asparaginy,

25 aspartyl,
 glutaminy,

30 D-homoseryl,
 3-(4-hydroxymethylphenyl)alanyl,
 isoleucyl,
 lysyl(N-epsilon-acetyl),
 methionyl,

35 3-(naphth-2-yl)alanyl,
 norvalyl,

octylglycyl,
prolyl,
3-(3-pyridyl)alanyl,
seryl,
5 D-seryl,
threonyl,
D-threonyl,
tryptyl,
tyrosyl, and
10 tyrosyl(O-methyl);

Xaa₇ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-glycyl, N-(R³)-isoleucyl, N-(R³)-leucyl, N-(R³)-D-leucyl, N-(R³)-norleucyl, N-(R³)-norvalyl, N-(R³)-seryl, N-(R³)-threonyl, and N-(R³)-valyl,
15 wherein R³ is as defined above; or Xaa₇ is an N-unalkylated amino acid selected from the group consisting of

alanyl,
allothreonyl,
allylglycyl,
20 3-(4-amidophenyl)alanyl,
2-aminobutyryl,
arginyl,
asparaginyll,
cyclohexylalanyl,
25 glutaminyll,
D-glutaminyll,
glycyl,
homoalanyl,
homoseryl,
30 4-hydroxyprolyl,
leucyl,
D-leucyl,
lysyl(N-epsilon-acetyl),
methionyl sulfone,
35 methionyl sulfoxide,
methionyl,

- norleucyl,
norvalyl,
D-norvalyl,
octylglycyl,
5 ornithyl(N-delta-acetyl),
phenylalanyl,
propargylglycyl,
seryl,
D-seryl,
10 threonyl,
tryptyl,
tyrosyl, and
valyl;
- 15 Xaa₈ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
alanyl, N-(R³)-D-alanyl, N-(R³)-isoleucyl, and N-(R³)-leucyl, wherein R³ is as
defined above; or Xaa₈ is an N-unalkylated amino acid selected from the group
consisting of
- 20 alanyl,
alloisoleucyl,
D-alloisoleucyl,
allylglycyl,
citrullyl,
glycyl,
25 isoleucyl,
D-isoleucyl,
leucyl,
D-leucyl,
lysyl(N-epsilon-acetyl),
30 D-lysyl(N-epsilon-acetyl),
methionyl,
3-(naphth-1-yl)alanyl,
norvalyl,
prolyl,
35 D-prolyl, and
valyl;

Xaa₉ is the N-alkylated amino acid N-(R³)-arginyl, wherein R³ is as defined above; or Xaa₉ is an N-unalkylated amino acid selected from the group consisting of

5 [(4-amino-N-isopropyl)cyclohexyl]alanyl,
 3-(4-amino-N-isopropylphenyl)alanyl,
 arginyl(N^GN^{G'} diethyl),
 arginyl,
 D-arginyl,
 citrullyl,
 10 glutaminyl,
 3-(4-guanidinophenyl)alanyl,
 histidyl,
 homoarginyl,
 lysyl(N-epsilon-isopropyl),
 15 lysyl(N-epsilon-nicotinyl),
 lysyl,
 norarginyl,
 ornithyl,
 ornithyl[N-delta-(2-imidazolinyl)],
 20 ornithyl(N-delta-isopropyl), and
 3-(3-pyridyl)alanyl;


25 Xaa₁₀ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-D-alanyl, N-(R³)-glycyl, N-(R³)-homoalanyl, and N-(R³)-norvalyl, wherein R³ is as defined above; or Xaa₁₀ is an N-unalkylated amino acid selected from the group consisting of

30 D-alanyl,
 2-aminobutyryl,
 D-2-aminobutyryl,
 2-aminoisobutyryl,
 3,4-dehydroprolyl,
 4-hydroxyprolyl,
 phenylalanyl,
 prolyl,
 35 D-prolyl,
 1,2,3,4-tetrahydroisoquinoline-3-carbonyl, and

D-valyl; and

Xaa₁₁ is a hydroxy group or an amino acid amide selected from the group consisting of:

- 5 alanylamine,
- D-alanylamine,
- alanylethylamine,
- D-alanylethylamine,
- azaglycylamine,
- 10 glycylamine,
- glycylethylamine,
- lysyl(N-epsilon-acetyl),
- D-lysyl(N-epsilon-acetyl),
- N-methyl-D-alanylamine,
- 15 sarcosylamine,
- serylamine,
- D-serylamine,
- a residue represented by the formula



- 20 a group represented by the formula $-\text{NH}-\text{R}^6$; wherein

s is an integer from 0 to 8;

R⁴ is selected from the group consisting of hydrogen, alkyl, and a 5- to 6-membered cycloalkyl ring;

- 25 R⁵ is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy;
- provided that s is not zero when R⁵ is hydroxy or alkoxy; and
- R⁶ is selected from hydrogen and hydroxy.

30 In another aspect, the present invention provides a composition for treating a patient in need of anti-angiogenesis therapy comprising a compound of formula (I) in combination with a pharmaceutically acceptable carrier.

 Yet another aspect of the present invention provides a method for treating a patient in need of anti-angiogenesis therapy comprising administering to the patient a therapeutically effective amount of a compound of formula (I).

Still yet another aspect of the present invention provides a composition for the treatment of a disease selected from cancer, arthritis, psoriasis, angiogenesis of the eye associated with infection or surgical intervention, macular degeneration, and diabetic retinopathy comprising a compound of formula (I) in combination with a pharmaceutically acceptable carrier.

In yet another aspect, the present invention provides a method of isolating a receptor from an endothelial cell comprising binding a compound of formula (I) to the receptor to form a peptide receptor complex, isolating the peptide receptor complex, and purifying the receptor.

Detailed Description of the Invention

Definition of Terms

As used herein, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise.

As used in the present specification the following terms have the meanings indicated:

The term "N-acetylamino," as used herein, refers to -NHC(O)CH_3 .

The term "acyl," as used herein, refers to an alkyl group attached to the parent molecular moiety through a carbonyl group.

The term "alkoxy," as used herein, refers to an alkyl group attached to the parent molecular moiety through an oxygen atom.

The term "alkyl," as used herein, refers to a monovalent group derived from a straight or branched chain saturated hydrocarbon by the removal of a hydrogen atom. Preferred alkyl groups for the present invention are alkyl groups having from one to five carbon atoms (C_1 - C_5 alkyl). Alkyl groups of one to three carbon atoms (C_1 - C_3 alkyl) are more preferred for the present invention.

The term "amino," as used herein, refers to -NH_2 .

The term "aryl," as used herein, represents a mono- or bicyclic carbocyclic ring system having one or two aromatic rings and is exemplified by phenyl, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, fluorenyl, indanyl, indenyl, and the like. The aryl groups of the present invention can be optionally substituted with one, two, three, four, or five substituents independently selected from the group consisting of alkoxy, alkyl, carboxy, and halo.

The term "carbonyl," as used herein, refers to -C(O)- .

The term "carboxy," as used herein, refers to CO_2H .

The term "cycloalkenyl," as used herein, refers to a non-aromatic cyclic or bicyclic

ring system having three to ten carbon atoms and one to three rings, wherein each five-membered ring has one double bond, each six-membered ring has one or two double bonds, each seven- and eight-membered ring has one to three double bonds, and each nine- to ten-membered ring has one to four double bonds. Examples of cycloalkenyl groups include cyclohexenyl, octahydronaphthalenyl, norbornylenyl, and the like. The cycloalkenyl groups of the present invention can be optionally substituted with one, two, three, four, or five substituents independently selected from the group consisting of alkoxy, alkyl, carboxy, halo, and hydroxy.

The term "cycloalkyl," as used herein, refers to a saturated monocyclic, bicyclic, or tricyclic hydrocarbon ring system having three to twelve carbon atoms. Examples of cycloalkyl groups include cyclopropyl, cyclopentyl, bicyclo[3.1.1]heptyl, adamantyl, and the like. The cycloalkyl groups of the present invention can be optionally substituted with one, two, three, four, or five substituents independently selected from the group consisting of alkoxy, alkyl, carboxy, halo, and hydroxy.

The term "halo," as used herein, refers to F, Cl, Br, or I.

The term "heterocycle," as used herein, refers to a five-, six-, or seven-membered ring containing one, two, or three heteroatoms independently selected from the group consisting of nitrogen, oxygen, and sulfur. The five-membered ring has zero to two double bonds and the six- and seven-membered rings have zero to three double bonds. The term "heterocycle" also includes bicyclic groups in which the heterocycle ring is fused to an aryl group. The heterocycle groups of the present invention can be attached through a carbon atom or a nitrogen atom in the group. Examples of heterocycles include, but are not limited to, furyl, thienyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, imidazoliny, pyrazolyl, isoxazolyl, isothiazolyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, pyridinyl, indolyl, indolinyl, benzothienyl, and the like. The heterocycle groups of the present invention can be optionally substituted with one, two, three, or four substituents independently selected from the group consisting of alkoxy, alkyl, carboxy, halo, and hydroxy.

The term "hydroxy," as used herein, refers to -OH.

The term "nicotinyl," as used herein, refers to the acyl group derived from nicotinic acid, i.e. pyridine-3-carboxylic acid. The term "2-Me-nicotinyl" or "2-methylnicotinyl" refers to a nicotinyl moiety substituted with a methyl group at the carbon adjacent to the nitrogen atom in the 2-position.

The term "nitrogen protecting group" or "N-protecting group," as used herein, refers to an easily removable group which is known in the art to protect an amino group against undesirable reaction during synthetic procedures and to be selectively removable.

The use of nitrogen protecting groups is well known in the art for protecting groups against undesirable reactions during a synthetic procedure and many such protecting groups are known (see, for example, T.H. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 2nd edition, John Wiley & Sons, New York (1991). Examples of N-protecting groups include, are not limited to, acyl groups including acetyl, trifluoroacetyl, acylisothiocyanate, aminocaproyl, benzoyl and the like, and acyloxy groups, including *t*-butyloxycarbonyl (Boc) and carbobenzyloxy (Cbz), 9-fluorenylmethoxycarbonyl (Fmoc), and the like.

The term "pharmaceutically acceptable ester," as used herein, refers to esters which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanolic, alkenolic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than six carbon atoms. Examples of particular esters include formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

The term "pharmaceutically acceptable prodrugs," as used herein, refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

The term "pharmaceutically acceptable salt," as used herein, represents salts or zwitterionic forms of the compounds of the present invention which are water or oil-soluble or dispersible, which are suitable for treatment of diseases without undue toxicity, irritation, and allergic response; which are commensurate with a reasonable benefit/risk ratio, and which are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting an amino group with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate,

formate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, 5 trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate, and undecanoate. Also, amino groups in the compounds of the present invention can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides. Examples of acids which can 10 be employed to form therapeutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric.

Basic addition salts can be prepared during the final isolation and purification of the compounds by reacting a carboxy group with a suitable base such as the hydroxide, 15 carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of pharmaceutically acceptable salts include lithium, sodium, potassium, calcium, magnesium, and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, 20 diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, 1-ephedrine, and N,N'-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

25 The term "pharmaceutically acceptable solvate," as used herein, refers to an aggregate that comprises one or more molecules of the solute, such as a compound of formula (I), with one or more molecules of solvent.

The term "receptor," as used herein, refers to chemical groups or molecules on the cell surface or in the cell interior that have an affinity for a specific chemical group or 30 molecule. Isolation of receptors relevant to the antiangiogenic activity of the peptide of the invention can provide useful diagnostic tools.

The term "shikimyl," as used herein, refers to the acyl residue derived from shikimic acid or [3R-(3 α ,4 α ,5 β)-3,4,5-trihydroxy]-1-cyclohexene-1-carboxylic acid. A "dihydroshikimyl" group denotes the fully saturated analog of shikimic acid.

35 The term "succinyl," as used herein, refers to the acyl residue derived from succinic acid or (1,4-dioxobutyl)-1-carboxylic acid.

Unless indicated otherwise by a "D-" prefix, *e.g.* D-Ala or N-Me-D-Ile, the stereochemistry of the α -carbon of the amino acids and aminoacyl residues in peptides described in this specification and the appended claims is the natural or "L" configuration. The Cahn-Ingold-Prelog "R" and "S" designations are used to specify the stereochemistry of chiral centers in certain acyl substituents at the N-terminus of the peptides of this invention. The designation "R,S" is meant to indicate a racemic mixture of the two enantiomeric forms. This nomenclature follows that described in R.S. Cahn, *et al.*, *Angew. Chem. Int. Ed. Engl.*, **5**, 385-415 (1966).

All peptide sequences are written according to the generally accepted convention whereby the α -N-terminal amino acid residue is on the left and the α -C-terminal is on the right. As used herein, the term " α -N-terminal" refers to the free α -amino group of an amino acid in a peptide, and the term " α -C-terminal" refers to the free α -carboxylic acid terminus of an amino acid in a peptide.

For the most part, the names on naturally occurring and non-naturally occurring aminoacyl residues used herein follow the naming conventions suggested by the IUPAC Commission on the Nomenclature of Organic Chemistry and the IUPAC-IUB Commission on Biochemical Nomenclature as set out in "Nomenclature of α -Amino Acids (Recommendations, 1974)" *Biochemistry*, **14**(2), (1975). To the extent that the names and abbreviations of amino acids and aminoacyl residues employed in this specification and appended claims differ from those suggestions, they will be made clear to the reader. Some abbreviations useful in describing the invention are defined below in the following Table 1.

Table 1

Abbreviation	Definition
N-Ac-Sar	N-acetylsarcosyl
Ala	alanyl
β -Ala	β -alanyl
AlaNH ₂	alanylamide
AlaNH-ethyl	alanylethylamide
alloIle	alloisoleucyl
alloThr	allothreonyl
alloThr(<i>t</i> -Bu)	allothreonyl(O- <i>t</i> -butyl)
Allylgly	allylglycyl
4-AmdPheAla	3-(4-amidophenyl)alanyl
2-Ambut	2-aminobutyryl

4-Ambut	4-aminobutyryl
(1R,3S)-AmCyCO	(1R,3S)-1-aminocyclopentane-3-carbonyl
(1S,3R)-AmCyCO	(1S,3R)-1-aminocyclopentane-3-carbonyl
(1R,4S)-AmCyeCO	(1R,4S)-1-aminocyclopent-2-ene-4-carbonyl
(1S,4R)-AmCyeCO	(1S,4R)-1-aminocyclopent-2-ene-4-carbonyl
2-Amisobut	2-aminoisobutyryl
4-AmIspCha	[(4-amino-N-isopropyl)cyclohexyl]alanyl
4-AmIspPheAla	3-(4-amino-N-isopropylphenyl)alanyl
4-AmPheAla	3-(4-aminophenyl)alanyl
Arg	arginyl
Arg(diethyl)	arginyl(N ^G N ^{G'} diethyl)
Asn	asparaginyl
Asn(Trt)	asparaginyl(trityl)
AzaGlyNH ₂	azaglycylamide
3-BzlThiAla	3-(3-benzothieryl)alanyl
5-BrThiAla	3-[2-(5-bromothieryl)]alanyl
Gly(<i>t</i> -Bu)	<i>t</i> -butylglycyl
3-ClPheAla	3-(3-chlorophenyl)alanyl
4-ClPheAla	3-(4-chlorophenyl)alanyl
Cit	citrullyl
3-CNPheAla	3-(3-cyanophenyl)alanyl
4-CNPheAla	3-(4-cyanophenyl)alanyl
Cha	cyclohexylalanyl
Chg	cyclohexylglycyl
Cys	cysteinyl
Cys(acme)	cysteinyl(S-acetamidomethyl)
Cys(<i>t</i> -Bu)	cysteinyl(S- <i>t</i> -butyl)
dePro	3,4-dehydropyrol
3,4-diFPheAla	3-(3,4-difluorophenyl)alanyl
3,4-diOMe-PheAla	3-(3,4-dimethoxyphenyl)alanyl
Fmoc	9-fluorenylmethyloxycarbonyl
3-FPheAla	3-(3-fluorophenyl)alanyl
4-FPheAla	3-(4-fluorophenyl)alanyl

Gln	glutaminyl
Gln(Trt)	glutaminyl(trityl)
GlyNH ₂	glycylamide
GlyNH-ethyl	glycylethylamide
4-GuPheAla	3-(4-guanidinophenyl)alanyl
Hala	homoalanyl
Harg	homoarginine
His	histidyl
HpheAla	homophenylalanyl
Hser	homoseryl
4-OHMePheAla	3-(4-hydroxymethylphenyl)alanyl
4-OHPro	4-hydroxypropyl
Ile	isoleucyl
Leu	leucyl
Lys	lysyl
Lys(Ac)	lysyl(N-epsilon-acetyl)
Lys(Isp)	lysyl(N-epsilon-isopropyl)
Lys(Nic)	lysyl(N-epsilon-nicotinyl)
Met	methionyl
Met(O ₂)	methionyl(sulfone)
Met(O)	methionyl(sulfoxide)
N-MeArg(Mtr)	(N ^G -4-methoxy-2,3,6-trimethylbenzenesulfonyl)arginyl
N-MeAla	N-methylalanyl
N-MeAlaNH ₂	N-methylalanyl amide
N-MeAlaNH-ethyl	N-methylalanyl ethylamide
N-MeAsp	N-methylaspartyl
N-MeAsp(<i>t</i> -Bu)	N-methylaspartyl(S- <i>t</i> -butyl)
N-MeGlu	N-methylglutamyl
N-MeGlu(<i>t</i> -Bu)	N-methylglutamyl(S- <i>t</i> -butyl)
N-MeIle	N-methylisoleucyl
N-MeLeu	N-methylleucyl
N-MeNle	N-methylnorleucyl
N-MeNva	N-methylnorvalyl
N-MeNvaNH-ethyl	N-methylnorvalyl ethylamide

4-MePheAla	3-(4-methylphenyl)alanyl
N-MePheAla	N-methylphenylalanyl
N-MePro	N-methylprolyl
N-MeSer(<i>t</i> -Bu)	N-methylseryl(O- <i>t</i> -butyl)
N-MeThr(Bzl)	N-methylthreonyl(O-benzyl)
N-MeThr(<i>t</i> -Bu)	N-methylthreonyl(O- <i>t</i> -butyl)
N-MeTyr	N-methyltyrosyl
N-MeTyr(<i>t</i> -Bu)	N-methyltyrosyl(O- <i>t</i> -butyl)
N-MeVal	N-methylvalyl
1-Nal	3-(naphth-1-yl)alanyl
2-Nal	3-(naphth-2-yl)alanyl
4-NO ₂ PheAla	3-(4-nitrophenyl)alanyl
Narg	norarginyl
Nle	norleucyl
Nor	norornithyl
Nva	norvalyl
octylgly	octylglycyl
Orn	ornithyl
Orn(Ac)	ornithyl(N-delta-acetyl)
Orn(Im)	ornithyl[N-delta-(2-imidazoliny)]
Orn(Isp)	ornithyl(N-delta-isopropyl)
Pen	penicillaminy
Pen(Sacme)	penicillaminy(S-acetamidomethyl)
Pen(SBzl)	penicillaminy(S-benzyl)
Pen(SMe)	penicillaminy(S-methyl)
3-PentaFPheAla	3-(pentafluorophenyl)alanyl
Arg(Pmc)	(N ^G -2,2,5,7,8-pentamethylchroman-6-sulfonyl)arginyl
PheAla	phenylalanyl
PheGly	phenylglycyl
Pro	prolyl
ProNH-ethyl	prolyl ethylamide
Propgly	propargylglycyl
3-PyrAla	3-(3-pyridyl)alanyl
Sar	sarcosyl

SarNH ₂	sarcosylamide
SarNH-ethyl	sarcosyl ethylamide
Ser	seryl
SerNH ₂	serylamide
Ser(Bzl)	seryl(O-benzyl)
Ser(<i>t</i> -Bu)	seryl(O- <i>t</i> -butyl)
StyAla	styrylalanyl
Tic	1,2,3,4-tetrahydroisoquinoline-3-carbonyl
4-ThzAla	3-(4-thiazolyl)alanyl
2-ThiAla	3-(2-thienyl)alanyl
Thr	threonyl
Thr(Bzl)	threonyl(O-benzyl)
Thr(<i>t</i> -Bu)	threonyl(O- <i>t</i> -butyl)
3-CF ₃ PheAla	3-(3-trifluoromethylphenyl)alanyl
3,4,5-TriFPheAla	3-(3,4,5-trifluorophenyl)alanyl
Tyr	tyrosyl
Tyr(<i>t</i> -Bu)	tyrosyl(O- <i>t</i> -butyl)
Tyr(Et)	tyrosyl(O-ethyl)
Tyr(Me)	tyrosyl(O-methyl)
Val	valyl

When not found in the table above, nomenclature and abbreviations may be further clarified by reference to the Calbiochem-Novabiochem Corp. *1999 Catalog and Peptide Synthesis Handbook* or the Chem-Impex International, Inc. *Tools for Peptide & Solid Phase Synthesis 1998-1999 Catalogue*.

In one aspect, the present invention relates to compounds of formula (I), wherein Xaa₂-Xaa₁₀ each represent an amino acyl residue; Xaa₁ may be absent or Xaa₁ is hydrogen, N-methylprolyl, or an acyl group; and Xaa₁₁ is a hydroxy group, an amino acid amide, or an amino residue. The amino acyl residues represented by Xaa₂-Xaa₁₀ can have an N-alkylated or an N-unalkylated amide bond. At least one of the amide bonds on a residue represented by Xaa₃, Xaa₄, Xaa₅, Xaa₆, Xaa₇, Xaa₈, Xaa₉, or Xaa₁₀ is N-alkylated.

Xaa₁ is absent or is selected from the group consisting of hydrogen; N-methylprolyl; R¹-(CH₂)_n-C(O)-, wherein n is an integer from 0 to 8 and R¹ is selected from the group consisting of N-acetylamino, alkoxy, alkyl, aryl, carboxy, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy; and R²-CH₂CH₂-O-(CH₂CH₂O)_p-CH₂-C(O)-,

wherein p is an integer from 1 to 8 and R² is selected from the group consisting of hydrogen, N-acetylamino, and alkyl. Preferably, Xaa₁ is absent or is selected from the group consisting of acetyl, N-methylprolyl, and succinyl.

Amino acyl residues suitable for the Xaa₂ position include N-methylalanyl, sarcosyl, N-ethylglycyl, N-methylnorvalyl, N-methylprolyl, β-alanyl, D-alanyl, 4-aminobutyryl, (1R,3S)-1-aminocyclopentane-3-carbonyl, (1S,3R)-1-aminocyclopentane-3-carbonyl, (1R,4S)-1-aminocyclopent-2-ene-4-carbonyl, (1S,4R)-1-aminocyclopent-2-ene-4-carbonyl, asparaginy, 3-(4-chlorophenyl)alanyl, 3-(4-cyanophenyl)alanyl, glutaminy, glutamyl, glycyl, 4-hydroxyprolyl, 3-(4-methylphenyl)alanyl, prolyl, seryl, and threonyl.

When Xaa₂ is an N-alkylated prolyl residue, Xaa₁ is absent. Preferably, Xaa₂ is an amino acyl residue selected from the group consisting of sarcosyl and N-methylprolyl.

Suitable amino acyl residues for Xaa₃ include N-methylalanyl, sarcosyl, N-methylleucyl, N-methylphenylalanyl, alanyl, (1S,3R)-1-aminocyclopentane-3-carbonyl, (1S,4R)-1-aminocyclopent-2-ene-4-carbonyl, asparaginy, aspartyl, 3-(3-cyanophenyl)alanyl, 3-(4-cyanophenyl)alanyl, glutaminy, glycyl, leucyl, lysyl(N-epsilon-acetyl), 3-(4-methylphenyl)alanyl, norvalyl, prolyl, and phenylalanyl. The preferred amino acid residues for Xaa₃ are N-methylalanyl and glycyl.

The N-alkylated residues suitable for Xaa₄ include N-methylalanyl, sarcosyl, N-methylhomophenylalanyl, N-methylisoleucyl, N-methylleucyl, N-methylnorvalyl, N-methylphenylalanyl, N-methyl-D-phenylalanyl, N-methylseryl, N-methyltyrosyl, N-methylvalyl, and N-methyl-D-valyl. N-Unalkylated amino acyl residues suitable for Xaa₄ are alanyl, alloisoleucyl, allylglycyl, 2-aminobutyryl, (1R,4S)-aminocyclopent-2-ene-4-carbonyl, asparaginy, aspartyl, 3-[2-(5-bromothieryl)]alanyl, 3-(3-chlorophenyl)alanyl, 3-(4-chlorophenylalanyl), 3-(3-cyanophenyl)alanyl, cyclohexylalanyl, 3-(3,4-dimethoxyphenyl)alanyl, 3-(3-fluorophenyl)alanyl, 3-(4-fluorophenylalanyl), glutaminy, glycyl, histidyl, homophenylalanyl, homoseryl, isoleucyl, leucyl, lysyl(N-epsilon-acetyl), methionyl, methionyl(sulfone), 3-(4-methylphenyl)alanyl, 3-(naphth-1-yl)alanyl, 3-(naphth-2-yl)alanyl, norornithyl, norvalyl, phenylalanyl, phenylglycyl, prolyl, 3-(3-pyridyl)alanyl, 3-(4-thiazolyl)alanyl, 3-(2-thienyl)alanyl, seryl, seryl(O-benzyl), styrylalanyl, tryptyl, tyrosyl, valyl, and D-valyl. Preferred amino acyl residues for Xaa₄ are N-methylalanyl, N-methylisoleucyl, N-methylleucyl, N-methylnorvalyl, N-methylphenylalanyl, N-methyl-D-phenylalanyl, N-methylvalyl, N-methyl-D-valyl, asparaginy, glutaminy, isoleucyl, phenylalanyl, and valyl.

The N-alkylated amino acyl residues suitable for Xaa₅ are N-methyl-D-homophenylalanyl, N-methyl-D-isoleucyl, N-methyl-D-leucyl, and N-(R³)-D-phenylalanyl. N-unalkylated amino acyl residues suitable for Xaa₅ are D-alanyl,

alloisoleucyl, D-alloisoleucyl, D-2-aminobutyryl, D-3-(4-aminophenyl)alanyl, D-asparaginyl, D-3-(3-benzothienyl)alanyl, D-*t*-butylglycyl, D-(4-chlorophenyl)alanyl, D-citrullyl, D-3-(3-cyanophenyl)alanyl, D-cyclohexylalanyl, cyclohexylglycyl, D-cysteiny(S-acetamidomethyl), D-cysteiny(S-*t*-butyl), D-3-(3,4-difluorophenyl)alanyl, D-
 5 (3,4-dimethoxyphenyl)alanyl, D-glutaminyl, glycyl, D-homophenylalanyl, D-homoseryl, isoleucyl, D-isoleucyl, D-leucyl, D-lysyl, D-lysyl(N-epsilon-nicotinyl), D-methionyl, D-3-(4-methylphenyl)alanyl, D-3-(naphth-1-yl)alanyl, D-3-(naphth-2-yl), D-3-(4-nitrophenyl)alanyl, D-norleucyl, D-ornithyl, D-penicillaminyl, D-penicillaminyl(S-acetamidomethyl), D-penicillaminyl(O-benzyl), D-penicillaminyl(S-methyl), D-3-
 10 (pentafluorophenyl)alanyl, D-phenylalanyl, D-prolyl, D-seryl, D-seryl(O-benzyl), D-(2-thienyl)alanyl, D-threonyl, D-threonyl(O-benzyl), D-3-(3-trifluoromethylphenyl)alanyl, D-(3,4,5-trifluorophenyl)alanyl, D-tryptyl, D-tyrosyl, D-tyrosyl(ethyl), and D-valyl. The amino acyl residues preferred for Xaa₅ are N-methyl-D-leucyl, D-alloisoleucyl, D-isoleucyl, D-leucyl, D-homophenylalanyl, and D-penicillaminyl(S-methyl).

15 Examples of N-alkylated amino acids suitable for Xaa₆ are N-methylaspartyl, N-methylglutamyl, sarcosyl, N-methylseryl, N-methylthreonyl, N-methylthreonyl(O-benzyl), and N-methyltyrosyl. N-unalkylated amino acyl residues suitable for Xaa₆ are alanyl, allothreonyl, D-allothreonyl, allylglycyl, glutaminyl, glycyl, histidyl, homoseryl, D-homoseryl, 3-(4-hydroxymethylphenyl)alanyl, isoleucyl, lysyl(N-epsilon-acetyl),
 20 methionyl, 3-(naphth-2-yl)alanyl, norvalyl, octylglycyl, prolyl, 3-(3-pyridyl)alanyl, seryl, D-seryl, threonyl, D-threonyl, tryptyl, tyrosyl, and tyrosyl(O-methyl). The preferred amino acyl residues for Xaa₆ are N-methylaspartyl, N-methylglutamyl, sarcosyl, N-methylseryl, N-methyltyrosyl, N-methylthreonyl(O-benzyl), allothreonyl, seryl, threonyl, and tyrosyl.

25 N-Alkylated amino acyl residues suitable for Xaa₇ are N-methylalanyl, sarcosyl, N-methylisoleucyl, N-methyleucyl, N-methyl-D-leucyl, N-methylnorleucyl, N-methylnorvalyl, N-methylseryl, N-methylthreonyl, and N-methylvalyl. The N-unalkylated amino acyl residues suitable for Xaa₇ are alanyl, allothreonyl, allylglycyl, 3-(4-amidophenyl)alanyl, 2-aminobutyryl, arginyl, asparaginyl, cyclohexylalanyl, glutaminyl,
 30 D-glutaminyl, glycyl, homoalanyl, homoseryl, 4-hydroxyprolyl, leucyl, D-leucyl, lysyl(N-epsilon-acetyl), methionyl, methionyl sulfone, methionyl sulfoxide, norleucyl, norvalyl, D-norvalyl, octylglycyl, ornithyl(N-delta-acetyl), phenylalanyl, propargylglycyl, seryl, D-seryl, threonyl, tryptyl, tyrosyl, and valyl. Preferably, the amino acyl residue for Xaa₇ is selected from the group consisting of N-methylalanyl, sarcosyl, N-methylisoleucyl, N-methyleucyl,
 35 N-methyl-D-leucyl, N-methylnorleucyl, N-methylnorvalyl, N-methylseryl, N-methylthreonyl, N-methylvalyl, norleucyl, norvalyl, and seryl.


Suitable N-alkyl amino acyl residues for Xaa₃ include N-methylalanyl, N-methyl-D-alanyl, N-methylisoleucyl, and N-methylleucyl. N-Unalkylated amino acyl residues suitable for Xaa₃ include alanyl, alloisoleucyl, D-alloisoleucyl, allylglycyl, citrullyl, glycyl, isoleucyl, D-isoleucyl, leucyl, D-leucyl, lysyl(N-epsilon-acetyl), D-lysyl(N-epsilon-acetyl), methionyl, 3-(naphth-1-yl)alanyl, norvalyl, prolyl, D-prolyl, and valyl. Preferably, the amino acyl residue for Xaa₃ is selected from the group consisting of N-methylalanyl, N-methyl-D-alanyl, N-methylisoleucyl, N-methylleucyl, isoleucyl, D-isoleucyl, and D-lysyl(N-epsilon-acetyl).

One of the N-alkylated amino acyl residues suitable for Xaa₃ is N-methylarginyl. N-Unalkylated amino acyl residues for Xaa₃ are selected from the group consisting of [(4-amino-N-isopropyl)cyclohexyl]alanyl, 3-(4-amino-N-isopropylphenyl)alanyl, arginyl, arginyl(N^GN^{G'}diethyl), citrullyl, glutaminyl, 3-(4-guanidinophenyl)alanyl, histidyl, homoarginyl, lysyl(N-epsilon-isopropyl), lysyl(N-epsilon-nicotinyl), lysyl, norarginyl, ornithyl, ornithyl(N-delta-imidazolyl), ornithyl(N-delta-isopropyl), and 3-(3-pyridyl)alanyl. Preferred amino acyl residues for Xaa₃ are arginyl and N-methylarginyl.

N-Alkylated amino acids suitable for the Xaa₁₀ position include N-methylalanyl, N-methyl-D-alanyl, sarcosyl, N-methylhomoalanyl, and N-methylnorvalyl. Other residues suitable for Xaa₁₀ include D-alanyl, 2-aminobutyryl, D-2-aminobutyryl, 2-aminoisobutyryl, 3,4-dehydroprolyl, 4-hydroxyprolyl, phenylalanyl, prolyl, D-prolyl, 1,2,3,4-tetrahydroisoquinoline-3-carbonyl, and D-valyl. The amino acyl residues preferred for Xaa₁₀ are N-methylalanyl, sarcosyl, N-methylnorvalyl, and prolyl.

Preferably, one or two residues selected from Xaa₃, Xaa₄, Xaa₅, Xaa₆, Xaa₇, Xaa₈, Xaa₉ and Xaa₁₀ has an N-alkylated amino acyl residue. The more preferred compounds of the invention have one N-alkylated amide bond on an amino acyl residue not including Xaa₁, as represented by Xaa₃, Xaa₄, Xaa₅, Xaa₆, Xaa₇, Xaa₈, Xaa₉ or Xaa₁₀.

Xaa₁₁ is a hydroxy group or an amino acid amide selected from the group consisting of alanylamine, D-alanylamine, alanylethylamine, D-alanylethylamine, azaglycylamine, glycylamine, glycylethylamine, lysyl(N-epsilon-acetyl), D-lysyl(N-epsilon-acetyl), N-methyl-D-alanylamine, sarcosylamine, serylamine, and D-serylamine; or Xaa₁₁ is a group represented by the formula



or a group represented by the formula -NH-R^6 , wherein s is an integer selected from 0 to 8; R⁴ is selected from hydrogen, alkyl, and a 5- to 6-membered cycloalkyl ring; R⁵ is selected from the group consisting of hydrogen, alkoxy, alkyl, aryl, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy, provided that s is not zero when R⁵ is hydroxy or

alkoxy; and R⁶ is selected from hydrogen and hydroxy. The preferred Xaa₁₁ groups for modifying the C-terminus of the invention are NH-ethyl and D-alanylamide.

Compositions

5 The compounds of the invention, including not limited to those specified in the examples, possess anti-angiogenic activity. As angiogenesis inhibitors, such compounds are useful in the treatment of both primary and metastatic solid tumors, including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney,
10 bladder and urothelium), female genital tract (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate, seminal vesicles, testes and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas (including those arising from bone and soft tissues as well as Kaposi's sarcoma)
15 and tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas). Such compounds may also be useful in treating solid tumors arising from hematopoietic malignancies such as leukemias (i.e. chloromas, plasmacytomas and the plaques and tumors of mycosis fungosides and cutaneous T-cell lymphoma/leukemia) as
20 as well as in the treatment of lymphomas (both Hodgkin's and non-Hodgkin's lymphomas). In addition, these compounds may be useful in the prevention of metastases from the tumors described above either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

 Further uses include the treatment and prophylaxis of autoimmune diseases such as
25 rheumatoid, immune and degenerative arthritis; various ocular diseases such as diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, retrolental fibroplasia, neovascular glaucoma, rubeosis, retinal neovascularization due to macular degeneration, hypoxia, angiogenesis in the eye associated with infection or surgical intervention, and other abnormal neovascularization conditions of the eye; skin diseases such as psoriasis;
30 blood vessel diseases such as hemangiomas, and capillary proliferation within atherosclerotic plaques; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation. Other uses include the treatment of diseases characterized by excessive or abnormal stimulation of endothelial cells, including not limited to intestinal adhesions,
35 Crohn's disease, atherosclerosis, scleroderma, and hypertrophic scars, i.e. keloids. Another use is as a birth control agent, by inhibiting ovulation and establishment of the

placenta. The compounds of the invention are also useful in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochela minutesalia quintosa*) and ulcers (*Helicobacter pylori*). The compounds of the invention are also useful to reduce bleeding by administration prior to surgery, especially for the
5 treatment of resectable tumors.

The compounds of the invention may be used in combination with other compositions and procedures for the treatment of diseases. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with a peptide of the present invention and then a peptide of the present invention may be subsequently
10 administered to the patient to extend the dormancy of micrometastases and to stabilize and inhibit the growth of any residual primary tumor. Additionally, the compounds of the invention may be combined with pharmaceutically acceptable excipients, and optionally sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions.

15 A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid-base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid),
20 polylactide co-glycolide (copolymers of lactic acid and glycolic acid) polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of
25 either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

When used in the above or other treatments, a therapeutically effective amount of one of the compounds of the present invention may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt form. By a "therapeutically effective
30 amount" of the compound of the invention is meant a sufficient amount of the compound to treat an angiogenic disease, (for example, to limit tumor growth or to slow or block tumor metastasis) at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of
35 sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated

and the severity of the disorder; activity of the specific compound employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or
5 coincidental with the specific compound employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Alternatively, a compound of the present invention may be administered as
10 pharmaceutical compositions containing the compound of interest in combination with one or more pharmaceutically acceptable excipients. A pharmaceutically acceptable carrier or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The compositions may be administered parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders,
15 ointments, drops or transdermal patch), rectally, or buccally. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions for parenteral injection comprise pharmaceutically-acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions,
20 as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper
25 fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of
30 microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

35 Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters),

poly(anhydrides), and (poly)glycols, such as PEG. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

- 5 The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

- 10 Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical administration, including those for inhalation, may be prepared as a dry powder which may be pressurized or non-pressurized. In non-pressurized powder compositions, the active ingredient in finely divided form may be used in admixture with a larger-sized pharmaceutically-acceptable inert carrier comprising particles having a size, for example, of up to 100 micrometers in diameter.
- 15 Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 micrometers.

- 20 Alternatively, the composition may be pressurized and contain a compressed gas, such as nitrogen or a liquified gas propellant. The liquified propellant medium and indeed the total composition is preferably such that the active ingredient does not dissolve therein to any substantial extent. The pressurized composition may also contain a surface active agent, such as a liquid or solid non-ionic surface active agent or may be a solid anionic surface active agent. It is preferred to use the solid anionic surface active agent in the form of a sodium salt.

- 25 A further form of topical administration is to the eye. A compound of the invention is delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the compound is maintained in contact with the ocular surface for a sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye, as for example the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous
- 30 humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically-acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an encapsulating material. Alternatively, the compounds of the invention may be injected directly into the vitreous and aqueous humour.

- 35 Compositions for rectal or vaginal administration are preferably suppositories which may be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository

wax which are solid at room temperature liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 *et seq.*

While the compounds of the invention can be administered as the sole active pharmaceutical agent, they may also be used in combination with one or more agents which are conventionally administered to patients for treating angiogenic diseases. For example, the compounds of the invention are effective over the short term to make tumors more sensitive to traditional cytotoxic therapies such as chemicals and radiation. The compounds of the invention also enhance the effectiveness of existing cytotoxic adjuvant anti-cancer therapies. The compounds of the invention may also be combined with other antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In particular, when used in the treatment of solid tumors, compounds of the invention may be administered with IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, angioinhibins, TNP-470, pentosan polysulfate, platelet factor 4, LM-609, SU-5416, CM-101, Tecogalan, plasminogen-K-5, vasostatin, vitaxin, vasculostatin, squalamine, marimastat or other MMP inhibitors, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, cisplatin, taxol, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, and the like as well as with radiation.

Total daily dose of the compositions of the invention to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily and more usually 1 to 300 mg/kg body weight.

It will be understood that agents which can be combined with the compound of the present invention for the inhibition, treatment or prophylaxis of angiogenic diseases are not limited to those listed above, include in principle any agents useful for the treatment or prophylaxis of angiogenic diseases.

- 5 The peptides of the invention may be used for the development of affinity columns for isolation of receptors relevant to the antiangiogenic activity of the peptide of the invention, e.g. TSP-1 receptor, in, for example, cultured endothelial cells. As is known in the art, isolation and purification of the receptor may be followed by amino acid sequencing to identify and isolate polynucleotides which encode the receptor.
- 10 Recombinant expression of this receptor would allow greater amounts of receptor to be produced, e.g. to produce a sufficient quantity for use in high throughput screening assays to identify other angiogenesis inhibitors.

Determination of Biological Activity

15 *In Vitro* Assay for Angiogenic Activity

The human microvascular endothelial (HMVEC) migration assay was run according to the procedure of S. S. Tolsma, O. V. Volpert, D. J. Good, W. F. Frazier, P. J. Polverini and N. Bouck, J. Cell Biol. 122, 497-511 (1993).

- 20 The HMVEC migration assay was carried out using Human Microvascular Endothelial Cells-Dermal (single donor) and Human Microvascular Endothelial Cells, (neonatal). The BCE or HMVEC cells were starved overnight in DME containing 0.1% bovine serum albumin (BSA). Cells were then harvested with trypsin and resuspended in DME with 0.1% BSA at a concentration of 1.5×10^6 cells per mL. Cells were added to the bottom of a 48 well modified Boyden chamber (Nucleopore Corporation, Cabin John, MD). The chamber was assembled and inverted, and cells were allowed to attach for 2
- 25 hours at 37 °C to polycarbonate chemotaxis membranes (5 µm pore size) that had been soaked in 0.1% gelatin overnight and dried. The chamber was then reinverted, and test substances (total volume of 50 µL), including activators, 15 ng/mL bFGF/VEGF, were added to the wells of the upper chamber. The apparatus was incubated for 4 hours at
- 30 37 °C. Membranes were recovered, fixed and stained (Diff Quick, Fisher Scientific) and the number of cells that had migrated to the upper chamber per 3 high power fields counted. Background migration to DME + 0.1 BSA was subtracted and the data reported as the number of cells migrated per 10 high power fields (400X) or, when results from multiple experiments were combined, as the percent inhibition of migration compared to a
- 35 positive control.

Representative compounds described in Examples 1 to 109 inhibited human endothelial cell migration in the above assay by at least 50% inhibition when tested at concentrations of 100 nM. Preferred compounds inhibited human endothelial cell migration by at least 51% when tested at concentrations of 10 nM, and more preferred compounds inhibited human endothelial cell migration by at least 51% at concentrations of 1 nM.

Synthesis of the Peptides

The polypeptides of the present invention may be synthesized by many techniques that are known to those skilled in the art. For solid phase peptide synthesis, a summary of the many techniques may be found in J.M. Stewart and J.D. Young, Solid Phase Peptide Synthesis, W.H. Freeman Co. (San Francisco), 1963 and J. Meienhofer, Hormonal Proteins and Peptides, vol. 2, p. 46, Academic Press (New York), 1973. For classical solution synthesis see G. Schroder and K. Lupke, The Peptides, vol. 1, Academic Press (New York), 1965.

Reagents, resins, amino acids, and amino acid derivatives are commercially available and can be purchased from Chem-Impex International, Inc. (Wood Dale, IL, U.S.A.) or Calbiochem-Novabiochem Corp. (San Diego, CA, U.S.A.) unless otherwise noted herein.

In general, these methods comprise the sequential addition of one or more amino acids or suitably protected amino acids to a growing peptide chain. Normally, either the amino or carboxy group of the first amino acid is protected by a suitable protecting group. The protected or derivatized amino acid can then be either attached to an inert solid support or utilized in solution by adding the next amino acid in the sequence having the complimentary (amino or carboxy) group suitably protected, under conditions suitable for forming the amide linkage. The protecting group is then removed from this newly added amino acid residue and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any remaining protecting groups (and any solid support) are removed sequentially or concurrently, to afford the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under conditions which do not racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide.

A particularly preferred method of preparing compounds of the present invention involves solid phase peptide synthesis.

In this particularly preferred method the α -amino function is protected by an acid or base sensitive group. Such protecting groups should have the properties of being stable to the conditions of peptide linkage formation, while being readily removable without destruction of the growing peptide chain or racemization of any of the chiral centers contained therein. Suitable protecting groups are 9-fluorenylmethyloxycarbonyl (Fmoc), *t*-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), biphenylisopropyl-oxycarbonyl, *t*-amyloxycarbonyl, isobornyloxycarbonyl, (α,α)-dimethyl-3,5-dimethoxybenzyloxycarbonyl, *O*-nitrophenylsulfenyl, 2-cyano-*t*-butyloxycarbonyl, and the like. The 9-fluorenylmethyloxycarbonyl (Fmoc) protecting group is preferred.

Particularly preferred side chain protecting groups are: for arginine and lysine: acetyl (Ac), adamantyloxycarbonyl, benzyloxycarbonyl (Cbz), *t*-butyloxycarbonyl (Boc), 4-methoxybenzenesulfonyl, N^G -4-methoxy-2,3,6-trimethylbenzenesulfonyl (Mtr), nitro, 2,2,5,7,8-pentamethylchroman-6-sulfonyl (Pmc), and *p*-toluenesulfonyl; for asparagine: trityl (Trt); for aspartyl: *t*-butyl (*t*-Bu); for glutamyl: *t*-butyl (*t*-Bu); for glutaminyl: trityl (Trt); for histidine: trityl (Trt), benzyl, benzyloxycarbonyl (Cbz), *p*-toluenesulfonyl and 2,4-dinitrophenyl; for penicillamine: methyl; for serine: *t*-butyl (*t*-Bu), benzyl and tetrahydropyranyl; for threonine: benzyl, and *t*-butyl (*t*-Bu); for tryptophan: formyl and *t*-butyloxycarbonyl (Boc); and for tyrosine: acetyl (Ac), benzyl, *O*-bromobenzyloxycarbonyl, *t*-butyl (*t*-Bu), cyclohexyl, cyclophenyl, 2,6-dichlorobenzyl, and isopropyl.

In the solid phase peptide synthesis method, the C-terminal amino acid is attached to a suitable solid support or resin. Suitable solid supports useful for the above synthesis are those materials which are inert to the reagents and reaction conditions of the stepwise condensation-deprotection reactions, as well as being insoluble in the media used. The preferred solid support for synthesis of C-terminal carboxy peptides is 4-hydroxymethylphenoxymethyl-copoly(styrene-1% divinylbenzene). The preferred solid support for C-terminal amide peptides is 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)phenoxy-acetamidoethyl resin available from Applied Biosystems.

The C-terminal amino acid is coupled to the resin by means of a coupling mediated by N,N' -dicyclohexylcarbodiimide (DCC), N,N' -diisopropylcarbodiimide (DIC), [*O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate] (HATU), or *O*-benzotriazol-1-yl- N,N,N',N' -tetramethyluroniumhexafluorophosphate (HBTU), with or without 4-dimethylaminopyridine (DMAP), 1-hydroxybenzotriazole (HOBT), benzotriazol-1-yloxy-tris(dimethylamino)phosphoniumhexafluorophosphate (BOP) or bis(2-oxo-3-oxazolidinyl)phosphine chloride (BOPCl), for about 1 to about 24 hours at a temperature of between 10 °C and 50 °C in a solvent such as dichloromethane or DMF.

When the solid support is 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamidoethyl resin, the Fmoc group is cleaved with a secondary amine, preferably piperidine, prior to coupling with the C-terminal amino acid as described above. The preferred reagents used in the coupling to the deprotected 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)phenoxyacetamidoethyl resin are *O*-benzotriazol-1-yl-N,N,N',N'-tetramethyluroniumhexafluorophosphate (HBTU, 1 equiv.) and 1-hydroxybenzotriazole (HOBT, 1 equiv.), or [*O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate] (HATU, 1 equiv.), in DMF.

The coupling of successive protected amino acids can be carried out in an automatic polypeptide synthesizer as is well known in the art. In a preferred embodiment, the α -amino function in the amino acids of the growing peptide chain are protected with Fmoc. The removal of the Fmoc protecting group from the N-terminal side of the growing peptide is accomplished by treatment with a secondary amine, preferably piperidine. Each protected amino acid is then introduced in about 3-fold molar excess and the coupling is preferably carried out in DMF. The coupling agent is normally *O*-benzotriazol-1-yl-N,N,N',N'-tetramethyluroniumhexafluorophosphate (HBTU, 1 equiv.) and 1-hydroxybenzotriazole (HOBT, 1 equiv.) or [*O*-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate] (HATU, 1 equiv.).

At the end of the solid phase synthesis, the polypeptide is removed from the resin and deprotected, either in succession or in a single operation. Removal of the polypeptide and deprotection can be accomplished in a single operation by treating the resin-bound polypeptide with a cleavage reagent, for example trifluoroacetic acid containing thianisole, water, or ethanedithiol.

In cases wherein the C-terminus of the polypeptide is an alkylamide, the resin is cleaved by aminolysis with an alkylamine. Alternatively, the peptide may be removed by transesterification, e.g. with methanol, followed by aminolysis or by direct transamidation. The protected peptide may be purified at this point or taken to the next step directly. The removal of the side chain protecting groups is accomplished using the cleavage cocktail described above.

The fully deprotected peptide is purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivitized polystyrene-divinylbenzene (for example, AMBERLITE® XAD); silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on SEPHADEX® G-25, LH-20 or countercurrent distribution; high

performance liquid chromatography (HPLC), especially reverse-phase HPLC on octyl- or octadecylsilyl-silica bonded phase column packing.

The foregoing may be better understood in light of the examples which are meant to describe compounds and process which can be carried out in accordance with the invention and are not intended as a limitation on the scope of the invention in any way.

Abbreviations which have been used the following examples are: NMP for N-methylpyrrolidinone; HATU for [O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate]; DMF for N,N-dimethylformamide, and TFA for trifluoroacetic acid.

EXAMPLE 1

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl

In the reaction vessel of an Applied Biosystems 433A peptide synthesizer was placed 0.1mM of Fmoc-Pro-Sieber ethylamide resin. Cartridges of 1mM amino acids were sequentially loaded. The Fastmoc 0.1 with previous peak monitoring protocol was used with the following synthetic cycle:

1. Resin solvated with NMP for about 5 minutes;
2. Resin washed with NMP for about 5 minutes;
3. Fmoc group removed using 50% piperidine solution in NMP for 5 minutes, resin washed, and sequence repeated 3 to 4 times;
4. Fmoc-amino acid activated with 1mM of 0.5M HATU in DMF;
5. Activated Fmoc-amino acid added to the reaction vessel followed by addition of 1mM of 2M diisopropylamine in NMP solution;
6. Fmoc-amino acid coupled for 20 minutes;
7. Resin washed and Fmoc-group removed using 50% piperidine in NMP.

The following protected amino acids were sequentially coupled to the resin using the above protocol:

Amino acid	Coupling time
1. Fmoc-Arg(Pmc)	20 minutes
2. Fmoc-Ile	20 minutes
3. Fmoc-NMeNva	20 minutes
4. Fmoc-Thr(<i>t</i> -Bu)	20 minutes
5. Fmoc-D-Ile	20 minutes
6. Fmoc-Val	20 minutes
7. Fmoc-Gly	20 minutes
8. Fmoc-Sar	20 minutes

9. acetic acid	20 minutes
----------------	------------

Upon completion of the synthesis the resin-bound peptide was washed with methanol three times and dried *in vacuo*, then treated with a (95:5) TFA/water solution (3 mL) at room temperature overnight. The resin was filtered and washed 3 times with methanol. The filtrates and the washes were combined and concentrated *in vacuo*. The residue was treated with diethyl ether and the precipitate was filtered to provide the crude peptide as an amorphous powder. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; $R_t = 4.259$ minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H); Amino Acid Anal.: 1.14 Pro; 1.70 Arg; 1.96 Ile; 0.47 Thr; 0.97 Val; 0.90 Gly; 0.98 Sar.

EXAMPLE 2

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-NMelle-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMelle for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-NMelle-Arg-ProNH-ethyl as a trifluoroacetate salt; $R_t = 4.416$ minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H); Amino Acid Anal.: 1.05 Pro; 1.26 Arg; 1.0 Ile; 0.54 Thr; 1.29 Nva; 1.02; Val; 0.94 Gly; 1.01 Sar.

EXAMPLE 3

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeAla-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeAla for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing

0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeAla-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 2.84 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 980 (M+H); Amino Acid Anal.: 0.99 Pro; 1.54 Arg; 2.10 Ile; 0.50 Thr; 0.95 Val; 0.96 Gly; 0.98 Sar.

EXAMPLE 4

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeVal for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.11 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H); Amino Acid Anal.: 1.03 Pro; 1.47 Arg; 2.06 Ile; 0.49 Thr; 0.96 Val; 1.01 Gly; 0.99 Sar.

EXAMPLE 5

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeAlaNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMeAla for Fmoc-Pro in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeAlaNH-ethyl as a trifluoroacetate salt; R_t = 2.84 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 982 (M+H); Amino Acid Anal.: 1.46 Arg; 2.02 Ile; 1.04 Nva; 0.47 Thr; 0.98 Val; 0.96 Gly; 1.03 Sar.

EXAMPLE 6

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-SarNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-Sar for Fmoc-Pro in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-SarNH-ethyl as a trifluoroacetate salt; R_t = 2.92 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 968 (M+H); Amino Acid Anal.: 1.49 Arg; 2.07 Ile; 1.05 Nva; 0.55 Thr; 0.98 Val; 0.96 Gly; 1.96 Sar.

EXAMPLE 7

N-Succinyl-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product was prepared by the procedure described in Example 1 with the following modification: coupling with acetic acid at the end of the synthesis was replaced with treatment of the peptide resin overnight with a 10-fold excess of succinic anhydride/pyridine in NMP. Upon completion of the synthesis, washing of the resin-bound peptide, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Succinyl-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 2.61 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1066 (M+H); Amino Acid Anal.: 1.59 Arg; 2.23 Ile; 0.50 Thr; 1.01 Val; 1.00 Gly; 1.02 Sar; 0.99 Pro.

EXAMPLE 8

N-Succinyl-Sar-Gly-Val-D-Leu-Thr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-Leu for Fmoc-D-Ile in Example 7. Upon completion of the synthesis, washing of the resin-bound peptide, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100%

acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-succinyl-Sar-Gly-Val-D-Leu-Thr-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 2.67 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1066 (M+H); Amino Acid Anal.: 1.61 Arg; 1.01 Ile; 1.10 Leu; 0.45 Thr; 0.95 Val; 0.95 Gly; 1.01 Sar; 0.93 Pro.

EXAMPLE 9

N-Ac-Sar-NMeAla-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeAla for Fmoc-Gly and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-NMeAla-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.43 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1022 (M+H).

EXAMPLE 10

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-NMeArg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMeArg(Mtr) for Fmoc-Arg(Pmc) in Example 1. Upon completion of the synthesis, cleavage of the peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-NMeArg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.43 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H).

EXAMPLE 11

N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeVal for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.42 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H).

EXAMPLE 12

N-Ac-Sar-Gly-NMeIle-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeIle for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMeIle-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.83 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1022 (M+H).

EXAMPLE 13

N-Ac-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMePheAla for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 4.00 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1056 (M+H).

EXAMPLE 14N-Ac-Sar-Gly-NMeNva-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeNva for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide from the resin, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column with a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMeNva-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; $R_t = 3.55$ minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1008 (M+H).

EXAMPLE 17N-Ac-Sar-Gly-NMeAla-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The procedure described in Example 1 was used substituting Fmoc-NMeAla for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMeAla-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) m/e 978 (M+H).

EXAMPLE 18N-MePro-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting NMePro for acetic acid and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-MePro-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) m/e 1063 (M+H).

EXAMPLE 21N-Ac-Sar-Gly-Val-D-Ile-NMeThr(Bzl)-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeThr(Bzl) for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-NMeThr(Bzl)-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) *m/e* 1084 (M+H).

EXAMPLE 22N-Ac-Sar-Gly-Val-D-Ile-Thr-Sar-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Sar for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-Sar-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; *R_t* = 2.68 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) *m/e* 966 (M+H); Amino Acid Anal.: 1.07 Pro; 1.21 Arg; 2.11 Ile; 0.47 Thr; 1.01; Val; 0.97 Gly; 2.07 Sar.

EXAMPLE 23N-Ac-Sar-Gly-Val-D-Leu-Sar-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-Leu for Fmoc-D-Ile, Fmoc-Sar for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Leu-Sar-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) *m/e* 964 (M+H).

EXAMPLE 24N-Ac-Sar-Gly-NMeLeu-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeLeu for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-NMeLeu-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) m/e 1022 (M+H).

EXAMPLE 25N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeLeu-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeLeu for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeLeu-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) m/e 1022 (M+H).

EXAMPLE 26N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; MS (ESI) m/e 1008 (M+H).

EXAMPLE 27N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeVal-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile and Fmoc-NMeVal for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeVal-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

10

EXAMPLE 28

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂

The desired product was prepared using the procedure in Example 1 with the following modifications: Fmoc-D-Ala-Sieber amide resin was substituted for Fmoc-Pro-Sieber ethylamide resin, and a coupling with Fmoc-Pro prior was added prior to the coupling with Fmoc-Arg(Pmc). Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂ as a trifluoroacetate salt; MS (ESI) m/e 1051 (M+H).

20

EXAMPLE 29

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂

The desired product was prepared by substituting Fmoc-NMeVal for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂ as a trifluoroacetate salt; R_t = 3.54 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1051 (M+H); Amino Acid Anal.: 1.00 Pro; 1.13 Arg; 2.07 Ile; 0.53 Thr; 1.05 Ala; 1.03 Val; 1.05 Gly; 1.03 Sar.

35

EXAMPLE 30N-Ac-Sar-Gly-Val-D-Ile-NMeThr-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-NMeThr(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can then be lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-NMeThr-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

EXAMPLE 31N-Ac-Sar-Gly-Val-D-Ile-NMeSer-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-NMeSer-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; $R_t = 3.01$ minutes (using a C-18 column and a solvent mixture increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 994.7 (M+H).

EXAMPLE 32N-Ac-Sar-Gly-Val-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-Leu for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

EXAMPLE 33

N-Ac-Sar-Gly-Val-D-Leu-Ser-NMeNva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-Leu for Fmoc-D-Ile and Fmoc-Ser(*t*-Bu) for Fmoc-Thr(*t*-Bu) in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-Leu-Ser-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

10

EXAMPLE 34

N-Ac-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-Ser(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 3.21 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 982 (M+H); Amino Acid Anal.: 1.02 Pro; 1.32 Arg; 2.12 Ile; 0.31 Ser; 1.01 Val; 1.03 Gly; 0.94 Sar.

25

EXAMPLE 35

N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile and Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 2.82 minutes (using a C-18 column and a

35

solvent mixture increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 996.7 (M+H).

EXAMPLE 36

5 N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a
10 solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt; R_t = 2.77 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI)
15 m/e 996.6 (M+H).

EXAMPLE 37

N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-DIle
20 and Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The
25 pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂ as a trifluoroacetate salt.

EXAMPLE 38

N-Ac-Sar-Gly-Phe-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by substituting Fmoc-PheAla for Fmoc-Val
30 and Fmoc-NMeVal for Fmoc-Nva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100%
35 acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions

can be lyophilized to provide N-Ac-Sar-Gly-Phe-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂ as a trifluoroacetate salt.

EXAMPLE 39

5 N-Ac-Sar-Gly-Val-D-allolle-Tyr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-allolle for Fmoc-D-Ile and Fmoc-Tyr(*t*-Bu) for Fmoc-Thr(*t*-Bu) in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative
10 HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-allolle-Tyr-NMeNva-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

15 EXAMPLE 40

N-Ac-Sar-Gly-Val-D-alloIle-Tyr-NMeVal-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-Tyr(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-NMeVal for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide,
20 removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-Tyr-NMeVal-Ile-Arg-ProNH-ethyl as a trifluoroacetate salt.

25

EXAMPLE 41

N-Ac-Sar-Gly-Gln-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by substituting Fmoc-Gln(Trt) for Fmoc-Val in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide,
30 removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Gln-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂ as a trifluoroacetate salt.

35

EXAMPLE 43

N-Ac-Sar-Gly-Val-D-alloIle-NMeThr-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeThr(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-NMeThr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

10

EXAMPLE 44

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt.

20

EXAMPLE 45

N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by substituting Fmoc-NMeVal for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt.

30

EXAMPLE 46

N-Ac-Sar-Gly-NMeVal-D-alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-NMeVal for Fmoc-Val, Fmoc-D-alloIle for Fmoc-D-Ile, and Fmoc-Nva for Fmoc NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting

35

groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-NMeVal-D-
5 alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

EXAMPLE 47

N-Ac-Sar-Gly-Val-D-HpheAla-Thr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-HpheAla for Fmoc-
10 D-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide
15 N-Ac-Sar-Gly-Val-D-HpheAla-Thr-NMeNva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

EXAMPLE 48

N-Ac-Sar-Gly-Val-D-HpheAla-Thr-NMeVal-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-HpheAla for Fmoc-
20 D-Ile and Fmoc-NMeVal for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The
25 pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-HpheAla-Thr-NMeVal-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

EXAMPLE 49

N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeNva-Ile-Arg-ProNH-ethyl

30 The desired product can be prepared by substituting Fmoc-D-Pen(SMe) for Fmoc-D-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing
35 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide

N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeNva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

EXAMPLE 50

5 N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeVal-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-Pen(SMe) for Fmoc-D-Ile and Fmoc-NMeVal for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeVal-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

15 EXAMPLE 51

N-Ac-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Ser(*t*-Bu) for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide N-Ac-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.21 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 982 (M+H); Amino Acid Anal.: 1.05 Pro; 0.93 Arg; 0.35 Ser; 2.01 Ile; 0.96 Val; 1.01 Gly; 0.99 Sar.

30 EXAMPLE 54

NAc-Sar-Gly-Val-NMe-D-Leu-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMe-D-Leu for Fmoc-D-Ile and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100%

acetonitrile/water containing 0.01% TFA over a period of 50 min. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-NMe-D-Leu-Thr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt, R_t = 3.87 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 min); MS (ESI) m/e 1008 (M+H); Amino Acid Anal.: 0.98 Pro; 0.99 Arg; 1.04 Ile; 0.99 Nva; 0.55 Thr; 1.06 Val; 0.98 Gly; 1.00 Sar.

EXAMPLE 55

NAc-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeNvaNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMeNva-Sieber ethylamide resin for Fmoc-Pro-Sieber ethylamide resin in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 min. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeNvaNH-ethyl as trifluoroacetate salt; R_t = 4.21 minutes (using a C-18 column and a solvent system increasing in gradient from 20% to 95% acetonitrile/water containing 10mM ammonium acetate over a period of 10 minutes); MS (ESI) m/e 1010 (M+H); Amino Acid Anal.: 0.99 Arg; 2.04 Ile; 1.03 Nva; 0.55 Thr; 1.0 Val; 0.97 Gly; 0.98 Sar.

EXAMPLE 56

NAc-Sar-Gly-Val-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMe-D-Leu for Fmoc-D-Ile, Fmoc-Ser(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.36 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 994.7 (M+H).

EXAMPLE 57NAc-Sar-Gly-Asn-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Asn(Trt) for Fmoc-Val, Fmoc-NMe-D-Leu for Fmoc-D-Ile, Fmoc-Ser(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Asn-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 2.39 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1009.7 (M+H).

EXAMPLE 58NAc-Sar-Gly-NMeNva-D-alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeNva for Fmoc-Val, Fmoc-D-alloIle for Fmoc-D-Ile, and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-NMeNva-D-alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.31 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1008.7 (M+H); Amino acid Anal.: 0.99 Pro; 0.99 Arg; 0.91 Nva; 0.49 Thr; 2.14 Ile; 0.97 Gly; 1.00 Sar.

EXAMPLE 59NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Asn(Trt) for Fmoc-Val, Fmoc-D-Leu for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water

containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; $R_t = 2.28$ minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1009.7 (M+H).

EXAMPLE 60

NAc-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂

The desired product was prepared by substituting Fmoc-NMePheAla for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt; $R_t = 3.45$ minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1099.7 (M+H).

EXAMPLE 61

NAc-Sar-Gly-Val-D-alloIle-NMeSer-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Val-D-alloIle-NMeSer-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

EXAMPLE 80

NAc-Sar-Gly-Val-D-Ile-Thr-NMeNle-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeNle for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a

solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-NMeNle-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.45 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1022 (M+H); Amino Acid Anal.: 1.03 Pro; 1.09 Arg; 0.45 Thr; 1.81 Ile; 1.07 Val; 1.01 Gly; 1.04 Sar.

EXAMPLE 81

NAc-Sar-Gly-Val-D-Ile-Sar-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Sar for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Sar-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.16 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 964.7 (M+H); Amino Acid Anal.: 1.01 Pro; 1.00 Arg; 0.92 Nva; 2.04 Ile; 1.04 Val; 0.99 Gly; 2.01 Sar.

EXAMPLE 82

NAc-Sar-Gly-Val-D-alloIle-Sar-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-Sar for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-alloIle-Sar-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.15 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 964.7 (M+H); Amino Acid Anal.: 1.00 Pro; 0.99 Arg; 0.90 Nva; 2.11 Ile; 1.03 Val; 0.96 Gly; 1.96 Sar.

EXAMPLE 83NAC-Sar-Gly-Val-D-Ile-Thr-Nva-NMeAla-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMeAla for Fmoc-Ile in Example 1. Upon completion of the synthesis,
5 cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMeAla-Arg-ProNH-ethyl
10 as trifluoroacetate salt; R_t = 2.86 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 966.7 (M+H); Amino Acid Anal.: 0.97 Pro; 0.98 Arg; 0.97 Nva; 0.51 Thr; 1.09 Ile; 1.02 Val; 0.96 Gly; 1.10 Sar.

15

EXAMPLE 84NAC-Sar-Gly-Val-D-Ile-NMeAsp-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeAsp(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and
20 precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-NMeAsp-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 2.89 minutes (using a C-18 column and a solvent
25 system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1022.7 (M+H); Amino Acid Anal.: 1.02 Pro; 1.02 Arg; 0.92 Nva; 2.01 Ile; 1.04 Val; 1.00 Gly; 0.98 Sar.

EXAMPLE 85

30

NAC-Sar-Gly-Val-D-Ile-Thr-NMe-D-Leu-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMe-D-Leu for NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent
35 system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-

Val-D-Ile-Thr-NMe-D-Leu-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.58 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1022.8 (M+H); Amino Acid Anal.: 1.04 Pro; 1.03 Arg; 0.47 Thr; 1.87 Ile; 1.06 Val; 1.01 Gly; 1.05 Sar.

EXAMPLE 86

NAc-Sar-Gly-Val-D-Ile-NMeGlu-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMeGlu(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-NMeGlu-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.12 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1036.7 (M+H); Amino Acid Anal.: 1.01 Pro; 1.0 Arg; 0.93 Nva; 2.04 Ile; 1.04 Val; 0.98 Gly; 1.0 Sar.

EXAMPLE 87

NAc-Sar-Gly-NMe-D-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMe-D-Val for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-NMe-D-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.12 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1008.7 (M+H); Amino Acid Anal.: 0.99 Pro; 1.02 Arg; 0.97 Nva; 0.43 Thr; 2.06 Ile; 0.96 Gly; 0.99 Sar.

EXAMPLE 88

NAc-Sar-Gly-Val-D-Ile-alloThr-NMeNle-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-*allo*Thr(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-NMeNle for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-*allo*Thr-NMeNle-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.59 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1022.8 (M+H); Amino Acid Anal.: 1.05 Pro; 0.97 Arg; 0.52 Thr; 1.88 Ile; 1.01 Gly; 1.03 Sar.

EXAMPLE 89

NAc-Sar-Gly-NMe-D-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-NMe-D-PheAla for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-NMe-D-Phe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.32 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1056.7 (M+H).

EXAMPLE 90

NAc-Sar-Gly-NMe-D-Phe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂

The desired product was prepared by substituting Fmoc-NMe-D-PheAla for Fmoc-Val and Fmoc-Nva for Fmoc-NMeNva in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-NMe-D-Phe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt; R_t = 3.18 minutes (using a C-18 column and a solvent

system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1099.7 (M+H).

EXAMPLE 91

5 NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMeLeu-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMeLeu for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative
10 HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMeLeu-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 3.39 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a
15 period of 10 minutes); MS (ESI) m/e 1008.7 (M+H).

EXAMPLE 92

NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Asn(Trt) for Fmoc-Val, Fmoc-D-Leu for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18
20 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 2.28 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a
25 period of 10 minutes); MS (ESI) m/e 1009.7 (M+H).

30

EXAMPLE 93

NAc-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-Pro-D-AlaNH₂

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-Ser(*t*-Bu) for Fmoc-Thr(*t*-Bu) and Fmoc-NMeSer(*t*-Bu) for Fmoc-NMeNva in
35 Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide

was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt; R_t = 2.74 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 1025.7 (M+H).

EXAMPLE 94

NAc-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeSer(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Ser(*t*-Bu) for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 2.53 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 982.6 (M+H).

EXAMPLE 95

NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMe-D-Ala-Arg-ProNH-ethyl

The desired product was prepared by substituting Fmoc-Nva for Fmoc-NMeNva and Fmoc-NMe-D-Ala for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide was obtained. This was purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions were lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMe-D-Ala-Arg-ProNH-ethyl as trifluoroacetate salt; R_t = 2.53 minutes (using a C-18 column and a solvent system increasing in gradient from 10% to 95% acetonitrile/water containing 0.01% TFA over a period of 10 minutes); MS (ESI) m/e 966.7 (M+H).

EXAMPLE 97

NAc-Sar-Gly-Val-D-alloIle-NMeTyr-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeTyr(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide,
5 removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Val-D-alloIle-NMeTyr-Nva-Ile-Arg-ProHEt as trifluoroacetate salt.

10

EXAMPLE 99NAc-Sar-Gly-Val-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-Ile for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide,
15 removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

20

EXAMPLE 100NAc-Sar-Gly-Val-D-Ile-alloThr-NMeNva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-alloThr(*t*-Bu) for Fmoc-Thr(*t*-Bu) in Example 1. Upon completion of the synthesis, cleavage of the resin-bound
25 peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Val-D-Ile-alloThr-NMeNva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

30

EXAMPLE 101NAc-Sar-Gly-Gln-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-Gln(Trt) for Fmoc-Val and Fmoc-D-Ile for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage
35 of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using

a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Gln-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

5

EXAMPLE 102

NAc-Sar-Gly-Val-D-Ile-Thr-NMeNva-D-Lys(Ac)-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-D-Lys(Ac) for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Val-D-Ile-Thr-NMeNva-D-Lys(Ac)-Arg-ProNH-ethyl as trifluoroacetate salt.

15

EXAMPLE 103

NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-Gln(Trt) for Fmoc-Val, Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeTyr(*t*-Bu) for Fmoc-Thr(*t*-Bu), and Fmoc-Nva for Fmoc-NMeNva in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

25

EXAMPLE 104

NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-D-Ile-Arg-ProNH-ethyl

The desired product can be prepared by substituting Fmoc-Gln(Trt) for Fmoc-Val, Fmoc-D-alloIle for Fmoc-D-Ile, Fmoc-NMeTyr(*t*-Bu) for Fmoc-Thr(*t*-Bu), Fmoc-Nva for Fmoc-NMeNva, and Fmoc-D-Ile for Fmoc-Ile in Example 1. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from

35

5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-D-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

5

EXAMPLE 105NAc-Sar-Gly-Phe-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂

The desired product can be prepared by substituting Fmoc-PheAla for Fmoc-Val in Example 28. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the protecting groups, and precipitation with diethyl ether, the crude peptide
10 can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to provide NAc-Sar-Gly-Phe-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂ as trifluoroacetate salt.

15

EXAMPLE 109NMePro-Gly-Ile-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl


The desired product can be prepared by substituting NMePro for Fmoc-Sar and Fmoc-Ile for Fmoc-Val in Example 1, and omitting the final coupling with acetic acid. Upon completion of the synthesis, cleavage of the resin-bound peptide, removal of the
20 protecting groups, and precipitation with diethyl ether, the crude peptide can be obtained. This can be purified by preparative HPLC using a C-18 column and a solvent system increasing in gradient from 5% to 100% acetonitrile/water containing 0.01% TFA over a period of 50 minutes. The pure fractions can be lyophilized to NMePro-Gly-Ile-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl as trifluoroacetate salt.

25

It will be evident to one skilled in the art that the instant invention is not limited to the forgoing illustrative examples, and that it can be embodied in other specific forms without departing from the essential attributes thereof. It is therefore desired that the examples be considered in all respects as illustrative and not restrictive, reference being
30 made to the appended claims, and all changes which come within the meaning and range of equivalency of the claims and therefore intended to be embraced therein.

WHAT IS CLAIMED IS:

1. A compound having a formula:



5

1 2 3 4 5 6 7 8 9

or a pharmaceutically acceptable salt thereof, wherein

at least one amide bond of an amino acid residue represented by Xaa₃, Xaa₄, Xaa₅, Xaa₆, Xaa₇, Xaa₈, Xaa₉, and Xaa₁₀ is N-alkylated;

10

Xaa₁ is absent or Xaa₁ is selected from the group consisting of hydrogen, N-methylpropyl, and an acyl group, wherein the acyl group is selected from the group consisting of

15

$\text{R}^1\text{-(CH}_2\text{)}_n\text{-C(O)-}$, wherein n is an integer from 0 to 8 and R¹ is selected from the group consisting of N-acetylamino, alkoxy, alkyl, aryl, carboxy, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy; and
 $\text{R}^2\text{-CH}_2\text{CH}_2\text{-O-(CH}_2\text{CH}_2\text{O)}_p\text{-CH}_2\text{-C(O)-}$, wherein p is an integer from 1 to 8 and R² is selected from the group consisting of hydrogen, N-acetylamino, and alkyl;

20

provided that Xaa₁ is absent only when Xaa₂ is N-(R³)-propyl;

25

Xaa₂ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-glycyl, N-(R³)-norvalyl, and N-(R³)-propyl, wherein R³ is C₁-C₅-alkyl; or Xaa₂ is an N-unalkylated amino acid selected from the group consisting of

30

-alanyl,
 D-alanyl,
 4-aminobutyryl,
 (1R,3S)-1-aminocyclopentane-3-carbonyl,
 (1S,3R)-1-aminocyclopentane-3-carbonyl,
 (1R,4S)-1-aminocyclopent-2-ene-4-carbonyl,
 (1S,4R)-1-aminocyclopent-2-ene-4-carbonyl,
 asparaginylyl,
 3-(4-chlorophenyl)alanyl,
 3-(4-cyanophenyl)alanyl,
 glutaminylyl,

35

- 40 glutamyl,
glycyl,
4-hydroxypropyl,
3-(4-methylphenyl)alanyl,
prolyl,
seryl, and
threonyl;
- 45 Xaa₃ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
alanyl, N-(R³)-glycyl, N-(R³)-leucyl, and N-(R³)-phenylalanyl, wherein R³ is as
defined above; or Xaa₃ is an N-unalkylated amino acid selected from the group
consisting of
alanyl,
50 (1S,3R)-1-aminocyclopentane-3-carbonyl,
(1S,4R)-1-aminocyclopent-2-ene-4-carbonyl,
asparaginyll,
aspartyl,
3-(3-cyanophenyl)alanyl,
55 3-(4-cyanophenyl)alanyl,
glutaminyll,
glycyl,
leucyl,
lysyl(N-epsilon-acetyl),
60 3-(4-methylphenyl)alanyl,
norvalyl,
prolyl, and
phenylalanyl;
- 65 Xaa₄ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
alanyl, N-(R³)-glycyl, N-(R³)-homophenylalanyl, N-(R³)-isoleucyl, N-(R³)-leucyl,
N-(R³)-norvalyl, N-(R³)-phenylalanyl, N-(R³)-D-phenylalanyl, N-(R³)-seryl, N-
(R³)-tyrosyl, N-(R³)-valyl, and N-(R³)-D-valyl, wherein R³ is as defined above; or
Xaa₄ is an N-unalkylated amino acid selected from the group consisting of
70 alanyl,
alloisoleucyl,
allylglycyl,

2-aminobutyryl,
(1R,4S)-aminocyclopent-2-ene-4-carbonyl,
75 asparaginyll,
aspartyl,
3-[2-(5-bromothieryl)]alanyl,
3-(3-chlorophenyl)alanyl,
3-(4-chlorophenyl)alanyl,
80 3-(3-cyanophenyl)alanyl,
cyclohexylalanyl,
3-(3,4-dimethoxyphenyl)alanyl,
3-(3-fluorophenyl)alanyl,
3-(4-fluorophenyl)alanyl,
85 glutaminyll,
glycyl,
histidyl,
homophenylalanyl,
homoseryl,
90 isoleucyl,
leucyl,
lysyl(N-epsilon-acetyl),
methionyl,
methionyl(sulfone),
95 3-(4-methylphenyl)alanyl,
3-(naphth-1-yl)alanyl,
3-(naphth-2-yl)alanyl,
norornithyl,
norvalyl,
100 phenylalanyl,
phenylglycyl,
prolyl,
3-(3-pyridyl)alanyl,
3-(4-thiazolyl)alanyl,
105 3-(2-thienyl)alanyl,
seryl,
seryl(O-benzyl),
styrylalanyl,

110 tryptyl,
 tyrosyl,
 valyl, and
 D-valyl;

115 Xaa₅ is an N-alkylated amino acid selected from the group consisting of N-(R³)-D-homophenylalanyl, N-(R³)-D-isoleucyl, N-(R³)-D-leucyl, and N-(R³)-D-phenylalanyl, wherein R³ is as defined above; or Xaa₅ is an N-unalkylated amino acid selected from the group consisting of

 D-alanyl,
 alloisoleucyl,
120 D-alloisoleucyl,
 D-2-aminobutyryl,
 D-3-(4-aminophenyl)alanyl,
 D-asparaginy,
 D-3-(3-benzothienyl)alanyl,
125 D-*t*-butylglycyl,
 D-(chlorophenyl)alanyl,
 D-citrullyl,
 D-3-(3-cyanophenyl)alanyl,
 D-cyclohexylalanyl,
130 cyclohexylglycyl,
 D-cysteinyl(S-acetamidomethyl),
 D-cysteinyl(S-*t*-butyl),
 D-3-(3,4-difluorophenyl)alanyl,
 D-(3,4-dimethoxyphenyl)alanyl,
135 D-glutaminy,
 glycyl,
 D-homophenylalanyl,
 D-homoseryl,
 isoleucyl,
140 D-isoleucyl,
 D-leucyl,
 D-lysyl(N-epsilon-nicotinyl),
 D-lysyl,
 D-methionyl,

- 145 D-3-(4-methylphenyl)alanyl,
D-3-(naphth-1-yl)alanyl,
D-3-(naphth-2-yl)alanyl,
D-3-(4-nitrophenyl)alanyl,
D-norleucyl,
- 150 D-ornithyl,
D-penicillaminyl(S-acetamidomethyl),
D-penicillaminyl(S-benzyl),
D-penicillaminyl(S-methyl),
D-penicillaminyl,
- 155 D-3-(pentafluorophenyl)alanyl,
D-phenylalanyl,
D-prolyl,
D-seryl(O-benzyl),
D-seryl,
- 160 D-(2-thienyl)alanyl,
D-threonyl(O-benzyl),
D-threonyl,
D-3-(3-trifluoromethylphenyl)alanyl,
D-(3,4,5-trifluorophenyl)alanyl,
- 165 D-tryptyl,
D-tyrosyl(O-ethyl),
D-tyrosyl, and
D-valyl;
- 170 Xaa₆ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
aspartyl, N-(R³)-glutamyl, N-(R³)-glycyl, N-(R³)-seryl, N-(R³)-threonyl, N-(R³)-
threonyl(O-benzyl), and N-(R³)-tyrosyl, wherein R³ is as defined above; or Xaa₆ is
an N-unalkylated amino acid selected from the group consisting of
alanyl,
- 175 allothreonyl,
D-allothreonyl,
allylglycyl,
asparaginyll,
aspartyl,
- 180 glutaminyll,

glyceryl,
 histidyl,
 homoseryl,
 D-homoseryl,
 185 3-(4-hydroxymethylphenyl)alanyl,
 isoleucyl,
 lysyl(N-epsilon-acetyl),
 methionyl,
 3-(naphth-2-yl)alanyl,
 190 norvalyl,
 octylglyceryl,
 prolyl,
 3-(3-pyridyl)alanyl,
 seryl,
 195 D-seryl,
 threonyl,
 D-threonyl,
 tryptyl,
 tyrosyl, and
 200 tyrosyl(O-methyl);

Xaa₇ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
 alanyl, N-(R³)-glyceryl, N-(R³)-isoleucyl, N-(R³)-leucyl, N-(R³)-D-leucyl, N-(R³)-
 norleucyl, N-(R³)-norvalyl, N-(R³)-seryl, N-(R³)-threonyl, and N-(R³)-valyl,
 205 wherein R³ is as defined above; or Xaa₇ is an N-unalkylated amino acid selected
 from the group consisting of

alanyl,
 allothreonyl,
 allylglyceryl,
 210 3-(4-amidophenyl)alanyl,
 2-aminobutyryl,
 arginyl,
 asparaginyl,
 cyclohexylalanyl,
 215 glutaminyl,
 D-glutaminyl,

glycyl,
homoalanyl,
homoseryl,
220 4-hydroxypropyl,
leucyl,
D-leucyl,
lysyl(N-epsilon-acetyl),
methionyl sulfone,
225 methionyl sulfoxide,
methionyl,
norleucyl,
norvalyl,
D-norvalyl,
230 octylglycyl,
ornithyl(N-delta-acetyl),
phenylalanyl,
propargylglycyl,
seryl,
235 D-seryl,
threonyl,
tryptyl,
tyrosyl, and
valyl;

240

Xaa₈ is an N-alkylated amino acid selected from the group consisting of N-(R³)-alanyl, N-(R³)-D-alanyl, N-(R³)-isoleucyl, and N-(R³)-leucyl, wherein R³ is as defined above; or Xaa₈ is an N-unalkylated amino acid selected from the group consisting of

245

alanyl,
alloisoleucyl,
D-alloisoleucyl,
allylglycyl,
citrullyl,
250 glycyl,
isoleucyl,
D-isoleucyl,

leucyl,
 D-leucyl,
 255 lysyl(N-epsilon-acetyl),
 D-lysyl(N-epsilon-acetyl),
 methionyl,
 3-(naphth-1-yl)alanyl,
 norvalyl,
 260 prolyl,
 D-prolyl, and
 valyl;

Xaa₉ is the N-alkylated amino acid N-(R³)-arginyl, wherein R³ is as defined above;
 265 or Xaa₉ is an N-unalkylated amino acid selected from the group consisting of

[(4-amino-N-isopropyl)cyclohexyl]alanyl,
 3-(4-amino-N-isopropylphenyl)alanyl,
 arginyl(N^GN^{G'} diethyl),
 arginyl,
 270 D-arginyl,
 citrullyl,
 glutaminyl,
 3-(4-guanidinophenyl)alanyl,
 histidyl,
 275 homoarginyl,
 lysyl(N-epsilon-isopropyl),
 lysyl(N-epsilon-nicotinyl),
 lysyl,
 norarginyl,
 280 ornithyl,
 ornithyl[N-delta-(2-imidazolinyl)],
 ornithyl(N-delta-isopropyl), and
 3-(3-pyridyl)alanyl;

285 Xaa₁₀ is an N-alkylated amino acid selected from the group consisting of N-(R³)-
 alanyl, N-(R³)-D-alanyl, N-(R³)-glycyl, N-(R³)-homoalanyl, and N-(R³)-norvalyl,
 wherein R³ is as defined above; or Xaa₁₀ is an N-unalkylated amino acid selected
 from the group consisting of


290 D-alanyl,
 2-aminobutyryl,
 D-2-aminobutyryl,
 2-aminoisobutyryl,
 3,4-dehydroprolyl,
 4-hydroxyprolyl,
 295 phenylalanyl,
 prolyl,
 D-prolyl,
 1,2,3,4-tetrahydroisoquinoline-3-carbonyl, and
 D-valyl; and

300

Xaa₁₁ is a hydroxy group or an amino acid amide selected from the group consisting of:

alanylamide,
 D-alanylamide,
 305 alanylethylamide,
 D-alanylethylamide,
 azaglycylamide,
 glycylamide,
 glycylethylamide,
 310 lysyl(N-epsilon-acetyl),
 D-lysyl(N-epsilon-acetyl),
 N-methyl-D-alanylamide,
 sarcosylamide,
 serylamide,
 315 D-serylamide,

a residue represented by the formula



a group represented by the formula $-\text{NH}-\text{R}^6$; wherein

s is an integer from 0 to 8;
 320 R⁴ is selected from the group consisting of hydrogen, alkyl, and a 5-
 to 6-membered cycloalkyl ring;
 R⁵ is selected from the group consisting of hydrogen, alkoxy, alkyl,
 aryl, cycloalkenyl, cycloalkyl, heterocycle, and hydroxy;

provided that s is not zero when R⁵ is hydroxy or alkoxy; and
325 R⁶ is selected from hydrogen and hydroxy.

2. A compound according to Claim 1, wherein Xaa₁ is absent or is selected from the group consisting of
- hydrogen,
 - acetyl,
 - 5 N-acetyl- alanyl,
 - butyryl,
 - (4-N-acetylamino)butyryl,
 - (6-N-acetylamino)caproyl,
 - (8-N-acetylamino)-3,6-dioxo-octanoyl,
 - 10 caproyl,
 - 5-chloro-2-hydroxynicotinyl,
 - 5-chloro-6-hydroxynicotinyl,
 - 2-chloroisonicotinyl,
 - 2-chloro-6-methylnicotinyl,
 - 15 cyclohexylacetyl,
 - furoyl,
 - 2-hydroxy-6-methylnicotinyl,
 - 6-hydroxynicotinyl,
 - 6-hydroxy-2-picolinyl,
 - 20 isonicotinyl,
 - 2-methoxyacetyl,
 - 2-methylnicotinyl,
 - 6-methylnicotinyl,
 - (4-methyl)phenylacetyl,
 - 25 N-methylprolyl,
 - nicotinyl,
 - phenylacetyl,
 - propionyl,
 - shikimyl,
 - 30 succinyl, and
 - tetrahydrofuroyl.

3. A compound according to Claim 2, wherein Xaa₁ is absent or is selected from the group consisting of

acetyl,
N-methylprolyl, and
succinyl.

4. A compound according to Claim 1, wherein Xaa₂ is selected from the group consisting of

N-methylalanyl,
sarcosyl,
N-ethylglycyl,
N-methylnorvalyl,
N-methylprolyl,
-alanyl,
4-aminobutyryl,
asparaginyll,
glutaminyll,
glutamyl,
glycyl,
prolyl,
seryl, and
threonyl.

5. A compound according to Claim 4, wherein Xaa₂ is selected from the group consisting of

sarcosyl, and
N-methylprolyl.

6. A compound according to Claim 1, wherein Xaa₃ is selected from the group consisting of

N-methylalanyl,
sarcosyl,
N-methyllaucyl,
N-methylphenylalanyl,
alanyl,
asparaginyll,

10 aspartyl,
 glutaminyll,
 glycyl,
 leucyl,
 norvalyl,
 prolyl, and
15 phenylalanyl.

7. A compound according to Claim 6, wherein Xaa₃ is selected from the group consisting of

 N-methylalanyl, and
 glycyl.

5

8. A compound according to Claim 1, wherein Xaa₄ is selected from the group consisting of

 N-methylalanyl,
 sarcosyl,
5 N-methylhomophenylalanyl,
 N-methylisoleucyl,
 N-methylleucyl,
 N-methylnorvalyl,
 N-methylphenylalanyl,
10 N-methyl-D-phenylalanyl,
 N-methylseryl,
 N-methyltyrosyl,
 N-methylvalyl,
 N-methyl-D-valyl,
15 3-[2-(5-bromothieryl)]alanyl,
 3-(3-chlorophenyl)alanyl,
 3-(4-chlorophenyl)alanyl,
 3-(3-cyanophenyl)alanyl,
 3-(3,4-dimethoxyphenyl)alanyl,
20 3-(3-fluorophenyl)alanyl,
 3-(4-fluorophenyl)alanyl,
 3-(4-methylphenyl)alanyl,
 3-(naphth-1-yl)alanyl,

25 3-(naphth-2-yl)alanyl,
3-(3-pyridyl)alanyl,
3-(4-thiazolyl)alanyl,
3-(2-thienyl)alanyl,
alloisoleucyl,
allylglycyl,
30 2-aminobutyryl,
asparaginy,
cyclohexylalanyl,
glutaminy,
glycyl,
35 histidyl,
homophenylalanyl,
homoseryl,
isoleucyl,
leucyl,
40 lysyl(N-epsilon-acetyl),
methionyl,
methionyl(sulfone),
norornithyl,
norvalyl,
45 phenylalanyl,
phenylglycyl,
prolyl,
seryl,
seryl(O-benzyl),
50 styrylalanyl,
tryptyl,
tyrosyl, and
valyl.

9. A compound according to Claim 8, wherein Xaa₄ is selected from the group consisting of

N-methylalanyl,
N-methylisoleucyl,
5 N-methylleucyl,

10 N-methylnorvalyl,
N-methylphenylalanyl,
N-methyl-D-phenylalanyl,
N-methylvalyl,
N-methyl-D-valyl,
asparaginyll,
glutaminyl,
isoleucyl,
phenylalanyl, and
15 valyl.

10. A compound according to Claim 1, wherein Xaa₅ is selected from the group consisting of

5 N-methyl-D-homophenylalanyl,
N-methyl-D-isoleucyl,
N-methyl-D-leucyl,
D-3-(4-aminophenyl)alanyl,
D-3-(3-benzothieryl)alanyl,
D-(chlorophenyl)alanyl,
D-3-(3-cyanophenyl)alanyl,
10 D-3-(3,4-difluorophenyl)alanyl,
D-(3,4-dimethoxyphenyl)alanyl,
D-3-(4-methylphenyl)alanyl,
D-3-(naphth-1-yl)alanyl,
D-3-(naphth-2-yl)alanyl.
15 D-3-(4-nitrophenyl)alanyl,
D-3-(pentafluorophenyl)alanyl,
D-3-(3-trifluoromethylphenyl)alanyl,
D-(3,4,5-trifluorophenyl)alanyl,
D-alanyl,
20 alloseucyl,
D-alloseucyl,
D-2-aminobutyryl,
D-asparaginyll,
D-citrullyl,
25 D-cyclohexylalanyl,

30 cyclohexylglycyl,
D-cysteinyl(S-acetamidomethyl),
D-cysteinyl(S-*t*-butyl),
D-glutaminy, glycy, 30
D-homophenylalanyl,
D-homoseryl,
isoleucyl,
D-isoleucyl,
35 D-leucyl,
D-lysyl(N-epsilon-nicotinyl),
D-lysyl,
D-methionyl,
D-norleucyl,
40 D-ornithyl,
D-penicillaminy(S-acetamidomethyl),
D-penicillaminy(S-benzyl),
D-penicillaminy(S-methyl),
D-penicillaminy,
45 D-phenylalanyl,
D-prolyl,
D-seryl(O-benzyl),
D-seryl,
D-*t*-butylglycyl,
50 D-(2-thienyl)alanyl,
D-threonyl(O-benzyl),
D-threonyl,
D-tryptyl,
D-tyrosyl(O-ethyl),
55 D-tyrosyl, and
D-valyl.

11. A compound according to Claim 10, wherein Xaa₅ is selected from the group consisting of

N-methyl-D-leucyl,
D-alloisoleucyl,

5 D-isoleucyl,
 D-leucyl,
 D-homophenylalanyl, and
 D-penacillaminy!(S-methyl).

12. A compound according to Claim 1, wherein Xaa₆ is selected from the group consisting of

 N-methylaspartyl,
 N-methylglutamyl,
5 sarcosyl,
 N-methylseryl,
 N-methyltyrosyl,
 N-methylthreonyl,
 N-methylthreonyl(O-benzyl),
10 alanyl,
 3-(4-hydroxymethylphenyl)alanyl,
 3-(naphth-2-yl)alanyl,
 3-(3-pyridyl)alanyl,
 allothreonyl,
15 D-allothreonyl,
 allylglycyl,
 glutaminy!,
 glycyl,
 histidyl,
20 homoseryl,
 D-homoseryl,
 isoleucyl,
 methionyl,
 norvalyl,
25 octylglycyl,
 prolyl,
 seryl,
 D-seryl,
 threonyl,
30 D-threonyl,
 tryptyl, and

tyrosyl.

13. A compound according to Claim 12, wherein Xaa₆ is selected from the group consisting of

5 N-methylaspartyl,
N-methylglutamyl,
sarcosyl,
N-methylseryl,
N-methyltyrosyl,
N-methylthreonyl,
N-methylthreonyl(O-benzyl),
10 allothreonyl,
seryl,
threonyl, and
tyrosyl.

14. A compound according to Claim 1, wherein Xaa₇ is selected from the group consisting of

5 N-methylalanyl,
sarcosyl,
N-methylisoleucyl,
N-methylleucyl,
N-methyl-D-leucyl,
N-methylnorleucyl,
N-methylnorvalyl,
10 N-methylseryl,
N-methylthreonyl,
N-methylvalyl,
alanyl,
allylglycyl,
15 3-(4-amidophenyl)alanyl,
2-aminobutyryl,
arginyl,
asparaginyll,
cyclohexylalanyl,
20 glutaminyll,

25 D-glutaminy,
glycyl,
homoalanyl,
homoseryl,
leucyl,
D-leucyl,
lysyl(N-epsilon-acetyl),
methionyl,
methionyl sulfone,
30 methionyl sulfoxide,
norleucyl,
norvalyl,
D-norvalyl,
octylglycyl,
35 ornithyl(N-delta-acetyl),
phenylalanyl,
propargylglycyl,
seryl,
D-seryl,
40 tyrosyl, and
valyl.

15. A compound according to Claim 14, wherein Xaa₇ is selected from the group consisting of

5 N-methylalanyl,
sarcosyl,
N-methylisoleucyl,
N-methylleucyl,
N-methyl-D-leucyl,
N-methylnorleucyl,
N-methylnorvalyl,
10 N-methylseryl,
N-methylthreonyl,
N-methylvalyl,
norleucyl,
norvalyl, and

15 seryl.

16. A compound according to Claim 1, wherein Xaa₈ is selected from the group consisting of

5 N-methylalanyl,
N-methyl-D-alanyl,
N-methylisoleucyl,
N-methyllleucyl,
3-(naphth-1-yl)alanyl,
alanyl,
allylglycyl,
10 glycyl,
isoleucyl,
D-isoleucyl,
leucyl,
D-lysyl(N-epsilon-acetyl),
15 methionyl,
norvalyl,
prolyl, and
valyl.

17. A compound according to Claim 16, wherein Xaa₈ is selected from the group consisting of

5 N-methylalanyl,
N-methyl-D-alanyl,
N-methylisoleucyl,
N-methyllleucyl,
isoleucyl,
D-isoleucyl, and
D-lysyl(N-epsilon-acetyl).

10 18. The compound according to Claim 1, wherein Xaa₉ is selected from the group consisting of

5 N-methylarginyl,
[(4-amino-N-isopropyl)cyclohexyl]alanyl,
3-(4-amino-N-isopropylphenyl)alanyl,

- 3-(4-guanidinophenyl)alanyl,
 arginyl,
 arginyl($N^G N^{G'}$ diethyl),
 citrullyl,
 10 2-[4-piperidinyl(N-amidino)]glycyl,
 glutaminyl,
 histidyl,
 homoarginyl,
 lysyl,
 15 lysyl(N-epsilon-isopropyl),
 lysyl(N-epsilon-nicotinyl),
 norarginyl,
 ornithyl,
 ornithyl[N-delta-(2-imidazoliny)], and
 20 ornithyl(N-delta-isopropyl).

19. A compound according to Claim 18, wherein Xaa₉ is selected from the group consisting of

arginyl, and
 N-methylarginyl.

5

20. A compound according to Claim 1, wherein Xaa₁₀ is selected from the group consisting of

N-methylalanyl,
 sarcosyl,
 5 N-methylhomoalanyl,
 N-methylnorvalyl,
 D-alanyl,
 2-aminobutyryl,
 2-aminoisobutyryl,
 10 3,4-dehydroprolyl,
 4-hydroxyprolyl,
 phenylalanyl,
 prolyl,
 D-prolyl, and

15 1,2,3,4-tetrahydroisoquinoline-3-carbonyl.

21. A compound according to Claim 20, wherein Xaa₁₀ is selected from the group consisting of

- 5 N-methylalanyl,
sarcosyl,
N-methylnorvalyl, and
prolyl.

22. A compound according to Claim 1, wherein Xaa₁₁ is selected from the group consisting of

- 5 alanylamine,
D-alanylamine,
alanylethylamine,
D-alanylethylamine,
azaglycylamine,
NH-cyclobutyl,
NH-cycloheptyl,
10 NH-1-(cyclohexyl)ethyl,
NH-2-(cyclohexyl)ethyl,
NH-2-(ethoxy)ethyl,
NH-ethyl,
NH-glycyl,
15 glycylethylamine,
NH-hexyl,
NH-2-(hydroxy)ethyl,
NH-isoamyl,
NH-isobutyl,
20 NH-2-(isopropoxy)ethyl,
NH-isopropyl,
lysyl(N-epsilon-acetyl),
D-lysyl(N-epsilon-acetyl),
NH-2-(methoxy)ethyl,
25 NH-3-(methoxy)propyl,
N-methyl-D-alanylamine,
NH-propyl,
NH-2-(1-pyrrolidine)ethyl,

30 sarcosylamide,
 serylamine, and
 D-serylamine.

23. A compound according to Claim 22, wherein Xaa₁ is selected from the group consisting of

 NH-ethyl, and
 D-alanylamine.

5

24. A compound according to Claim 1, wherein

 Xaa₁ is absent or is selected from the group consisting of
 acetyl,
5 N-methylprolyl, and
 succinyl;

 Xaa₂ is selected from the group consisting of
 sarcosyl, and
10 N-methylprolyl;

 Xaa₃ is selected from the group consisting of
 N-methylalanyl, and
 glycyl;

15

 Xaa₄ is selected from the group consisting of
 N-methylalanyl,
 N-methylisoleucyl,
 N-methylleucyl,
20 N-methylnorvalyl,
 N-methylphenylalanyl,
 N-methyl-D-phenylalanyl,
 N-methylvalyl,
 N-methyl-D-valyl,
25 asparaginyll,
 glutaminyll,
 isoleucyl,

phenylalanyl, and
valyl;

30

Xaa₅ is selected from the group consisting of

N-methyl-D-leucyl,
D-alloisoleucyl,
D-isoleucyl,
D-leucyl,
D-homophenylalanyl, and
D-penacillaminy(S-methyl);

35

Xaa₆ is selected from the group consisting of

N-methylaspartyl,
N-methylglutamyl,
sarcosyl,
N-methylseryl,
N-methyltyrosyl,
N-methylthreonyl,
N-methylthreonyl(O-benzyl),
allothreonyl,
seryl,
threonyl, and
tyrosyl;

40

45

50

Xaa₇ is selected from the group consisting of

N-methylalanyl,
sarcosyl,
N-methylisoleucyl,
N-methylleucyl,
N-methyl-D-leucyl,
N-methylnorleucyl,
N-methylnorvalyl,
N-methylseryl,
N-methylthreonyl,
N-methylvalyl,
norleucyl,

55

60

65 norvalyl, and
seryl;

70 Xaa₈ is selected from the group consisting of
N-methylalanyl,
N-methyl-D-alanyl,
N-methylisoleucyl,
N-methylleucyl,
isoleucyl,
D-isoleucyl, and
D-lysyl(N-epsilon-acetyl);

75 Xaa₉ is selected from the group consisting of
arginyl, and
N-methylarginyl;

80 Xaa₁₀ is selected from the group consisting of
N-methylalanyl,
sarcosyl,
N-methylnorvalyl, and
prolyl; and

85 Xaa₁₁ is selected from the group consisting of
NH-ethyl, and
D-alanylamide.

25. A compound according to Claim 24 wherein Xaa₁ is selected from the group consisting of

acetyl, and
succinyl.

5

26. A compound according to Claim 24 wherein Xaa₂ is sarcosyl.

27. A compound according to Claim 24 wherein Xaa₄ is selected from the group consisting of

N-methylleucyl,

5 N-methylnorvalyl,
N-methylphenylalanyl,
N-methyl-D-phenylalanyl, and
valyl.

28. A compound according to Claim 24 wherein Xaa₅ is selected from the group consisting of

5 N-methyl-D-leucyl,
D-alloisoleucyl,
D-isoleucyl, and
D-leucyl;

29. A compound according to Claim 24 wherein Xaa₆ is selected from the group consisting of

5 sarcosyl,
N-methylseryl,
N-methyltyrosyl,
allothreonyl,
seryl, and
threonyl.

30. A compound according to Claim 24 wherein Xaa₇ is selected from the group consisting of

5 N-methylalanyl,
N-methylnorvalyl,
N-methylvalyl, and
norvalyl.

31. A compound according to Claim 24 wherein Xaa₈ is selected from the group consisting of

5 N-methyllleucyl, and
isoleucyl.

32. A compound according to Claim 24 wherein Xaa₉ is arginyl.

33. A compound according to Claim 24 wherein Xaa₁₀ is selected from the group consisting of

N-methylalanyl, and
prolyl.

5

34. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

35. A method of treating a patient in need of anti-angiogenesis therapy comprising administering to the patient in need a therapeutically effective amount of a compound of Claim 1.

36. A composition for the treatment of a disease selected from cancer, arthritis, psoriasis, angiogenesis of the eye associated with infection or surgical intervention, macular degeneration and diabetic retinopathy comprising a compound of Claim 1 in combination with a pharmaceutically acceptable carrier.

5

37. A method of isolating a receptor from an endothelial cell comprising binding compound of Claim 1 to the receptor to form a peptide receptor complex; isolating the peptide receptor complex; and purifying the receptor.

38. A compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of

N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-SarNH-ethyl,
N-Succinyl-Sar-Gly-Val-D-Leu-Thr-NMeNva-Ile-Arg-ProNH-ethyl,

5 N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-NMeArg-ProNH-ethyl,

N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-NMeIle-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-NMeAla-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,

N-MePro-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,

10 N-Ac-Sar-Gly-Val-D-Ile-NMeThr(Bzl)-Nva-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-Val-D-Ile-Thr-Sar-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeLeu-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeVal-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂,

15 N-Ac-Sar-Gly-Val-D-Ile-NMeThr-Nva-Ile-Arg-ProNH-ethyl,

- N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Phe-D-Ile-Thr-NMeVal-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Val-D-alloIle-Tyr-NMeNva-Ile-Arg-ProNH-ethyl,
N-Ac-Sar-Gly-Val-D-alloIle-Tyr-NMeVal-Ile-Arg-ProNH-ethyl,
20 N-Ac-Sar-Gly-Gln-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-Val-D-alloIle-NMeThr-Nva-Ile-Arg-ProNH-ethyl,
N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-NMeVal-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
N-Ac-Sar-Gly-NMeVal-D-alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl,
25 N-Ac-Sar-Gly-Val-D-Hphe-Thr-NMeNva-Ile-Arg-ProNH-ethyl,
N-Ac-Sar-Gly-Val-D-Hphe-Thr-NMeVal-Ile-Arg-ProNH-ethyl,
N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeNva-Ile-Arg-ProNH-ethyl,
N-Ac-Sar-Gly-Val-D-Pen(SMe)-Thr-NMeVal-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeNvaNH-ethyl,
30 NAc-Sar-Gly-Val-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Asn-NMe-D-Leu-Ser-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-alloIle-NMeSer-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-NMeNle-Ile-Arg-ProNH-ethyl,
35 NAc-Sar-Gly-Val-D-Ile-Sar-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-alloIle-Sar-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMeAla-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-NMeAsp-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-NMe-D-Leu-Ile-Arg-ProNH-ethyl,
40 NAc-Sar-Gly-Val-D-Ile-NMeGlu-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-NMe-D-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-NMe-D-Phe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMeLeu-Arg-ProNH-ethyl,
NAc-Sar-Gly-Asn-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl,
45 NAc-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-Nva-NMe-D-Ala-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Val-D-Ile-alloThr-NMeNva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Gln-D-Ile-Thr-NMeNva-D-Ile-Arg-ProNH-ethyl,
50 NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-Ile-Arg-ProNH-ethyl,
NAc-Sar-Gly-Gln-D-alloIle-NMeTyr-Nva-D-Ile-Arg-ProNH-ethyl,

N-Ac-Sar-Gly-Phe-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂, and
NMePro-Gly-Ile-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl.

39. A compound or a pharmaceutically acceptable salt thereof, selected from the group consisting of
- N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-NMeIle-Arg-ProNH-ethyl,
 - 5 N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeAla-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeVal-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Ile-Thr-Nva-Ile-Arg-NMeAlaNH-ethyl,
 - N-Succinyl-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-NMeAla-Val-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - 10 N-Ac-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-NMeNva-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Leu-Sar-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-NMeLeu-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeNva-Ile-Arg-ProNH-ethyl,
 - 15 N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-Ile-Arg-Pro-D-AlaNH₂,
 - N-Ac-Sar-Gly-Val-D-Ile-NMeSer-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Leu-NMeSer-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Leu-Ser-NMeNva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-ProNH-ethyl,
 - 20 N-Ac-Sar-Gly-Val-D-alloIle-Thr-NMeSer-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeSer-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-alloIle-NMeSer-Ser-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-NMe-D-Leu-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-NMeNva-D-alloIle-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - 25 N-Ac-Sar-Gly-NMePhe-D-Ile-Thr-Nva-Ile-Arg-Pro-D-AlaNH₂,
 - N-Ac-Sar-Gly-Val-D-Ile-alloThr-NMeNle-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-NMe-DPhe-D-Ile-Thr-Nva-Ile-Arg-ProNH-ethyl,
 - N-Ac-Sar-Gly-Val-D-alloIle-Ser-NMeSer-Ile-Arg-Pro-D-AlaNH₂,
 - N-Ac-Sar-Gly-Val-D-alloIle-NMeTyr-Nva-Ile-Arg-ProNH-ethyl, and
 - 30 N-Ac-Sar-Gly-Val-D-Ile-Thr-NMeNva-DLys(Ac)-Arg-ProNH-ethyl.

SEQUENCE LISTING

<110> Abbott Laboratories
Haviv, Fortuna
Henkin, Jack
Kalvin, Douglas M.
Bradley, Michael F.

<120> N-ALKYLATED PEPTIDES HAVING
ANTIANGIOGENIC ACTIVITY

<130> 6632.PC.O1

<140> Not Yet Assigned
<141>

<150> Unknown
<151> 2000-10-31

<150> US 09/447,099
<151> 1999-11-22

<160> 1

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Antiangiogenetic Peptide

<221> VARIANT
<222> (1)...(1)
<223> Xaa = N-methylprolyl at position 1

<221> VARIANT
<222> (2)...(2)
<223> Xaa = N-(R3)Ala, N-(R3)Gly, N-(R3)Nva, N-(R3)Pro,
B-Ala, Asn, 4-ClPheAla, 4-CNPheAla, Gln, Glu, Gly,
4-OHPro, 4-MePheAla, Pro, Ser, or Thr at position
2

<221> VARIANT
<222> (3)...(3)
<223> Xaa = N-(R3)Ala, N-(R3)Gly, N-(R3)Leu,
N-(R3)PheAla, Ala, Asn, Asp, 3-CNPheAla,
4-CNPheAla, Gln, Gly, Leu, Lys(Ac), 4-MePheAla,
Nva, Pro, and PheAla as position 3

<221> VARIANT
<222> (4)...(4)
<223> Xaa = N-(R3)Ala, N-(R3)Gly, N-(R3)HpheAla,
N-(R3)Ile, N-(R3)Leu, N-(R3)Nva, N-(R3)PheAla,
N-(R3)Ser, N-(R3)Tyr, N-(R3)Val, Ala, AlloIle,

Allylgly, 2-Ambut, Asn, Asp at position 4

<221> VARIANT

<222> (4)...(4)

<223> Xaa = 5-BrThiAla, 3-ClPheAla, 4-ClPheAla,
3-CNPheAla, Cha, 3,4-diOMe-PheAla, 3-FpheAla,
4-FpheAla, Gln, Gly, His, HpheAla, Hser, Ile, Leu,
Lys(Ac), Met, Met(O2), 4-MePheAla at position 4

<221> VARIANT

<222> (4)...(4)

<223> Xaa = 1-Nal, 2-Nal, Nor, Nva, PheAla, PheGly,
Pro, 3-PyrAla, 4-ThzAla, 2-ThiAla, Ser, Ser(Bzl),
StyAla, Trp, Tyr, Val at position 4

<221> VARIANT

<222> (5)...(5)

<223> Xaa = AlloIle, Chg, Gly, Ile at position 5

<221> VARIANT

<222> (6)...(6)

<223> Xaa = N-(R3)Asp, N-(R3)Glu, N-(R3)Gly, N-(R3)Ser,
N-(R3)Thr, N-(R3)Thr(Bzl), N-(R3)Tyr, Ala,
AlloThr, Allylgly, Asn, Asp, Gln, Gly, His, Hser,
4-OHMePheAla, Ile, Lys(Ac) at position 6

<221> VARIANT

<222> (6)...(6)

<223> Xaa = Met, 2-Nal, Nva, Octylgly, Pro, 3-PyrAla,
Ser, Thr, Trp, Tyr, Tyr(Me) at position 6

<221> VARIANT

<222> (7)...(7)

<223> Xaa = N-(R3)Ala, N-(R3)Gly, N-(R3)Ile, N-(R3)Leu,
N-(R3)Nle, N-(R3)Nva, N-(R3)Ser, N-(R3)Thr,
N-(R3)Val, Ala, AlloThr, Allylgly, 4-AmdPheAla,
2-Ambut, Arg, Asn at position 7

<221> VARIANT

<222> (7)...(7)

<223> Xaa = Cha, Gln, Gly, Hala, Hser, 4-OHPro, Leu,
Lys(Ac), Met(O2), Met(O), Met, Nle, Nva,
Octylgly, Orn(Isp), PheAla, ProGly, Ser, Thr, Trp,
Tyr, and Val at position 7

<221> VARIANT

<222> (8)...(8)

<223> N-(R3)Ala, N-(R3)Ile, N-(R3)Leu, Ala, AlloIle,
Allylgly, Cit, Gly, Ile, Leu, Lys(Ac), Met, 1-Nal,
Nva, Pro, and Val at position 8

<221> VARIANT

<222> (9)...(9)

<223> Xaa = N-(R3)Arg, 4-AmIspCha, 4-AmIspPheAla,
Arg(diethyl), Arg, Cit, Gln, 4-GuPheAla, His,
Harg, Lys(Isp), Lys(Nic), Lys, Nor, Orn, Orn(Im),
Orn(Isp), and 3-PyrAla at position 9

<221> VARIANT

<222> (10)...(10)

<223> Xaa = N-(R3)Ala, N-(R3)Gly, N-(R3)Hala, N-(R3)Nva,
2-Ambut, 2-Amisobut, dePro, 4-OHPro, PheAla, Pro,
and Tic at position 10

<221> VARIANT

<222> (11)...(11)

<223> Xaa = AlanH2, AlanH-ethyl, AzaGlyNH2, GlyNH2,
GlyNH-ethyl, Lys(Ac), SarNH2, and SerNH2 at
position 11

<400> 1

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10

INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 00/32105

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/78 C07K7/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 41824 A (ABBOTT LAB) 13 November 1997 (1997-11-13) claim 1	1-39
Y	WO 98 41542 A (UNIV NORTHWESTERN ; ABBOTT LAB (US)) 24 September 1998 (1998-09-24) claim 1	1-39
Y	WO 97 33174 A (TORREY PINES INST FOR MOLECULA) 12 September 1997 (1997-09-12) claim 1	1-39



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

12 March 2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Deffner, C-A

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/US 00/32105

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9741824 A	13-11-1997	US 5801146 A	01-09-1998
		US 5981484 A	09-11-1999
		AU 724077 B	14-09-2000
		AU 3060697 A	26-11-1997
		BR 9708911 A	03-08-1999
		CZ 9803426 A	12-05-1999
		EP 0910571 A	28-04-1999
		US 5972896 A	26-10-1999
		US 6057122 A	02-05-2000
WO 9841542 A	24-09-1998	EP 0918795 A	02-06-1999
WO 9733174 A	12-09-1997	AU 720632 B	08-06-2000
		AU 2040597 A	22-09-1997
		CA 2248078 A	12-09-1997
		EP 0890101 A	13-01-1999